

=> d his

(FILE 'HOME' ENTERED AT 12:47:52 ON 14 JAN 2005)

FILE 'HCAPLUS' ENTERED AT 12:47:58 ON 14 JAN 2005

L1 1 (US20040082637 OR US6667302 OR US20020151575)/PN
 E WO1998-US27822/AP, PRN
 L2 1 WO1998-US27822/AP, PRN
 E US1997-070287/AP, PRN
 L3 1 US1997-070287P/AP, PRN
 L4 1 L1-3

FILE 'REGISTRY' ENTERED AT 12:50:28 ON 14 JAN 2005

FILE 'HCAPLUS' ENTERED AT 12:50:31 ON 14 JAN 2005

L5 TRA L4 1- RN : 9 TERMS

FILE 'REGISTRY' ENTERED AT 12:50:31 ON 14 JAN 2005

L6 9 SEA L5

FILE 'WPIX' ENTERED AT 12:50:34 ON 14 JAN 2005

L7 1 (US20040082637 OR US6667302 OR US20020151575)/PN
 L8 1 WO1998-US27822/AP, PRN
 L9 1 US1997-070287P/AP, PRN
 L10 1 L7-9

=> b hcap

FILE 'HCAPLUS' ENTERED AT 12:51:44 ON 14 JAN 2005

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 14 Jan 2005 VOL 142 ISS 3

FILE LAST UPDATED: 12 Jan 2005 (20050112/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d all 14

L4 ANSWER 1 OF 1 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1999:460416 HCAPLUS

DN 131:87914

ED Entered STN: 28 Jul 1999

TI Heterocyclic topoisomerase poisons, namely 2-(benzimidazol-5-yl)benzimidazoles

IN Lavoie, Edmond J.; Kim, Jun Sung; Liu, Leroy Fong

PA Rutgers, the State University of New Jersey, USA

SO PCT Int. Appl., 29 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM C07D403-04

ICS A61K031-415

CC 28-9 (Heterocyclic Compounds (More Than One Hetero Atom))

Section cross-reference(s): 1, 7

FAN.CNT 1


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9933824	A1	19990708	WO 1998-US27822	19981230 <--
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,				

Search done by Noble Jarrell

TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU,
 TJ, TM
 RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES,
 FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI,
 CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
 CA 2316221 AA 19990708 CA 1998-2316221 19981230 <--
 AU 9920220 A1 19990719 AU 1999-20220 19981230 <--
 AU 753268 B2 20021010
 EP 1044199 A1 20001018 EP 1998-965021 19981230 <--
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
 IE, FI
 JP 2002509858 T2 20020402 JP 2000-526506 19981230 <--
 US 2002151575 A1 20021017 US 2001-869141 20010613 <--
 US 6667302 B2 20031223
 US 2004082637 A1 20040429 US 2003-690800 20031021 <--
 PRAI US 1997-70287P P 19971231 <--
 WO 1998-US27822 W 19981230 <--
 US 2001-869141 A3 20010613

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
WO 9933824	ICM	C07D403-04
	ICS	A61K031-415
WO 9933824	ECLA	C07D235/20; C07D403/14+239+235C+235C; C07D403/14+249+235C+235C
US 2002151575	ECLA	C07D235/20; C07D403/14+239+235C+235C; C07D403/14+249+235C+235C
US 2004082637	ECLA	C07D235/20; C07D403/14+239+235C+235C; C07D403/14+249+235C+235C
OS MARPAT 131:87914		
GI		



AB The invention provides title compds. I [R₁, R₂ = H, alkyl, cycloalkyl, alkoxy, (un)substituted (hetero)aryl, etc; or R₁R₂ = benzo, methylenedioxy; R₃ = H, alkyl, cycloalkyl, alkoxy, OH, CF₃O, halo, etc.; R₄R₅ = 3- to 5-atom ring-forming chain containing .gtoreq.1 NH group, and as further units O (except peroxides), S, N(X), C, or C(O), where X = null, H, O, alkyl, Ph, or PhCH₂], as well as their pharmaceutically acceptable salts, pharmaceutical compns., and use of any of these to treat cancer. For instance, 5-phenyl-2-[2-(3,4-dinitrophenyl)benzimidazol-5-yl]benzimidazole was hydrogenated over Pd/C to give the 3,4-diamino compound, which underwent diazotization with concomitant cyclization to give title compound II. Two example compds. potently inhibited topoisomerase I in vitro, and also exhibited cytotoxic activity against RPMI 8402 cancer cells and camptothecin-resistant CPT-K5 cells in vitro.

ST heterocyclic topoisomerase poison benzimidazolylbenzimidazole prepn

IT Antitumor agents

Cytotoxic agents

(preparation of (benzimidazolyl)benzimidazoles as topoisomerase poisons for use as anticancer agents)

IT Antitumor agents

(solid tumor, treatment; preparation of (benzimidazolyl)benzimidazoles as topoisomerase poisons for use as anticancer agents)

IT 167959-27-5, 5-Phenyl-2-[2-(benzimidazol-5-yl)benzimidazol-5-yl]benzimidazole
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)
(comparison compound; preparation of (benzimidazolyl)benzimidazoles as topoisomerase poisons for use as anticancer agents)

IT 230308-98-2P, 5-Phenyl-2-[2-(quinoxalin-6-yl)benzimidazol-5-yl]benzimidazole
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
(comparison compound; preparation of (benzimidazolyl)benzimidazoles as topoisomerase poisons for use as anticancer agents)

IT 230308-95-9P, 5-Phenyl-2-[2-(3,4-diaminophenyl)benzimidazol-5-yl]benzimidazole
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(intermediate; preparation of (benzimidazolyl)benzimidazoles as topoisomerase poisons for use as anticancer agents)

IT 143180-75-0
RL: BPR (Biological process); BSU (Biological study, unclassified); MSC (Miscellaneous); BIOL (Biological study); PROC (Process)
(preparation of (benzimidazolyl)benzimidazoles as topoisomerase poisons for use as anticancer agents)

IT 144-62-7, Ethanedioic acid, reactions 517-21-5, Glyoxal disodium bisulfite 192879-72-4, 5-Phenyl-2-[2-(3,4-dinitrophenyl)benzimidazol-5-yl]benzimidazole
RL: RCT (Reactant); RACT (Reactant or reagent)
(starting material; preparation of (benzimidazolyl)benzimidazoles as topoisomerase poisons for use as anticancer agents)

IT 230308-96-0P, 5-Phenyl-2-[2-(1H-benzotriazol-5-yl)benzimidazol-5-yl]benzimidazole 230308-97-1P, 5-Phenyl-2-[2-(1,2,3,4-tetrahydro-2,3-dioxoquinoxalin-6-yl)benzimidazol-5-yl]benzimidazole
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(target compound; preparation of (benzimidazolyl)benzimidazoles as topoisomerase poisons for use as anticancer agents)

RE.CNT 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Arznad; ARZNEIM-FORSCH 1974, V24(12), P1927
- (2) Goldman, G; BIOCHEMISTRY 1997, V36(21), P6488 HCAPLUS
- (3) Heinz, L; US 3538097 A 1970
- (4) Kim, J; J MED CHEM 1996, V39(4), P992 HCAPLUS
- (5) Kim, J; JOURNAL OF MEDICINAL CHEMISTRY 1997, V40(18), P2818 HCAPLUS
- (6) Lavoie, E; WO 9636612 A 1996 HCAPLUS
- (7) Lavoie, E; WO 9831673 A 1998 HCAPLUS
- (8) Loewe, H; Basic substituted 2,6-bisbenzimidazole derivatives, a novel series of substances with chemotherapeutic activity 1975, 17, HCAPLUS
- (9) Sun, Q; BIOORGANIC & MEDICINAL CHEMISTRY LETTERS 1994, V4(24), P2871 HCAPLUS

=> b reg
FILE 'REGISTRY' ENTERED AT 12:51:51 ON 14 JAN 2005
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2005 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 12 JAN 2005 HIGHEST RN 812631-13-3
DICTIONARY FILE UPDATES: 12 JAN 2005 HIGHEST RN 812631-13-3

TSCA INFORMATION NOW CURRENT THROUGH MAY 21, 2004

Please note that search-term pricing does apply when conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.


Experimental and calculated property data are now available. For more information enter HELP PROP at an arrow prompt in the file or refer

Search done by Noble Jarrell

to the file summary sheet on the web at:
<http://www.cas.org/ONLINE/DBSS/registryss.html>

=> d ide l6 tot


L6 ANSWER 1 OF 9 REGISTRY COPYRIGHT 2005 ACS on STN
 RN 230308-98-2 REGISTRY
 CN Quinoxaline, 6-(5-phenyl[2,5'-bi-1H-benzimidazol]-2'-yl)- (9CI) (CA INDEX NAME)
 OTHER NAMES:
 CN 5-Phenyl-2-[2-(quinoxalin-6-yl)benzimidazol-5-yl]benzimidazole
 FS 3D CONCORD
 MF C28 H18 N6
 SR CA
 LC STN Files: CA, CAPLUS, TOXCENTER, USPAT2, USPATFULL
 DT.CA Caplus document type: Journal; Patent
 RL.P Roles from patents: BIOL (Biological study); PREP (Preparation)
 RL.NP Roles from non-patents: BIOL (Biological study); PREP (Preparation); PRP (Properties); USES (Uses)



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

3 REFERENCES IN FILE CA (1907 TO DATE)
 3 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L6 ANSWER 2 OF 9 REGISTRY COPYRIGHT 2005 ACS on STN
 RN 230308-97-1 REGISTRY
 CN 2,3-Quinoxalinedione, 1,4-dihydro-6-(5-phenyl[2,5'-bi-1H-benzimidazol]-2'-yl)- (9CI) (CA INDEX NAME)
 OTHER NAMES:
 CN 5-Phenyl-2-[2-(1,2,3,4-tetrahydro-2,3-dioxoquinoxalin-6-yl)benzimidazol-5-yl]benzimidazole
 FS 3D CONCORD
 MF C28 H18 N6 O2
 SR CA
 LC STN Files: CA, CAPLUS, TOXCENTER, USPAT2, USPATFULL
 DT.CA Caplus document type: Journal; Patent
 RL.P Roles from patents: BIOL (Biological study); PREP (Preparation); USES (Uses)
 RL.NP Roles from non-patents: BIOL (Biological study); PREP (Preparation); PRP (Properties); USES (Uses)




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

3 REFERENCES IN FILE CA (1907 TO DATE)
 3 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L6 ANSWER 3 OF 9 REGISTRY COPYRIGHT 2005 ACS on STN
 RN 230308-96-0 REGISTRY
 CN 1H-Benzotriazole, 5-(5-phenyl[2,5'-bi-1H-benzimidazol]-2'-yl)- (9CI) (CA INDEX NAME)
 OTHER NAMES:
 CN 5-Phenyl-2-[2-(1H-benzotriazol-5-yl)benzimidazol-5-yl]benzimidazole
 FS 3D CONCORD
 MF C26 H17 N7

Search done by Noble Jarrell

SR CA
 LC STN Files: CA, CAPLUS, TOXCENTER, USPAT2, USPATFULL
 DT.CA Caplus document type: Journal; Patent
 RL.P Roles from patents: BIOL (Biological study); PREP (Preparation); USES (Uses)
 RL.NP Roles from non-patents: BIOL (Biological study); PREP (Preparation); PRP (Properties); USES (Uses)




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

3 REFERENCES IN FILE CA (1907 TO DATE)
 3 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L6 ANSWER 4 OF 9 REGISTRY COPYRIGHT 2005 ACS on STN
 RN 230308-95-9 REGISTRY
 CN 1,2-Benzenediamine, 4-(5-phenyl[2,5'-bi-1H-benzimidazol]-2'-yl)- (9CI)
 (CA INDEX NAME)

OTHER NAMES:

CN 5-Phenyl-2-[2-(3,4-diaminophenyl)benzimidazol-5-yl]benzimidazole
 FS 3D CONCORD
 MF C26 H20 N6
 SR CA
 LC STN Files: CA, CAPLUS, TOXCENTER, USPAT2, USPATFULL
 DT.CA Caplus document type: Journal; Patent
 RL.P Roles from patents: PREP (Preparation); RACT (Reactant or reagent)
 RL.NP Roles from non-patents: PREP (Preparation); RACT (Reactant or reagent)




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

4 REFERENCES IN FILE CA (1907 TO DATE)
 4 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L6 ANSWER 5 OF 9 REGISTRY COPYRIGHT 2005 ACS on STN
 RN 192879-72-4 REGISTRY
 CN 2,5'-Bi-1H-benzimidazole, 2'-(3,4-dinitrophenyl)-5-phenyl- (9CI) (CA INDEX NAME)

OTHER NAMES:


CN 5-Phenyl-2-[2-(3,4-dinitrophenyl)benzimidazol-5-yl]benzimidazole
 FS 3D CONCORD
 MF C26 H16 N6 O4
 SR CA
 LC STN Files: CA, CAPLUS, TOXCENTER, USPAT2, USPATFULL
 DT.CA Caplus document type: Journal; Patent
 RL.P Roles from patents: RACT (Reactant or reagent)
 RL.NP Roles from non-patents: PREP (Preparation); RACT (Reactant or reagent)



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

4 REFERENCES IN FILE CA (1907 TO DATE)
4 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L6 ANSWER 6 OF 9 REGISTRY COPYRIGHT 2005 ACS on STN
RN 167959-27-5 REGISTRY
CN 2,5':2',5''-Ter-1H-benzimidazole, 5-phenyl- (9CI) (CA INDEX NAME)
OTHER NAMES:
CN 5-Phenyl-2-[2-(benzimidazol-5-yl)benzimidazol-5-yl]benzimidazole
FS 3D CONCORD
MF C27 H18 N6
SR CA
LC STN Files: CA, CAPLUS, TOXCENTER, USPAT2, USPATFULL
DT.CA Caplus document type: Journal; Patent
RL.P Roles from patents: BIOL (Biological study); PREP (Preparation); USES (Uses)
RL.NP Roles from non-patents: BIOL (Biological study); PREP (Preparation); PROC (Process); PRP (Properties); USES (Uses)



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

16 REFERENCES IN FILE CA (1907 TO DATE)
16 REFERENCES IN FILE CAPLUS (1907 TO DATE)


L6 ANSWER 7 OF 9 REGISTRY COPYRIGHT 2005 ACS on STN
RN 143180-75-0 REGISTRY
CN Isomerase, deoxyribonucleate topo-, I (9CI) (CA INDEX NAME)
OTHER NAMES:
CN Deoxyribonucleate topoisomerase I
CN DNA topoisomerase I
CN Topiosomerase I
MF Unspecified
CI MAN
SR CA
LC STN Files: AGRICOLA, BIOBUSINESS, BIOSIS, CA, CAPLUS, CEN, CHEMCATS, CIN, IPA, PROMT, TOXCENTER, USPAT2, USPATFULL
DT.CA Caplus document type: Book; Conference; Dissertation; Journal; Patent
RL.P Roles from patents: ANST (Analytical study); BIOL (Biological study); FORM (Formation, nonpreparative); MSC (Miscellaneous); OCCU (Occurrence); PREP (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses)
RLD.P Roles for non-specific derivatives from patents: BIOL (Biological study); PREP (Preparation); PROC (Process); PRP (Properties); USES (Uses)
RL.NP Roles from non-patents: ANST (Analytical study); BIOL (Biological study); FORM (Formation, nonpreparative); MSC (Miscellaneous); OCCU (Occurrence); PREP (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses)
RLD.NP Roles for non-specific derivatives from non-patents: ANST (Analytical study); BIOL (Biological study); FORM (Formation, nonpreparative); OCCU (Occurrence); PREP (Preparation); PROC (Process); PRP (Properties); USES (Uses)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

2318 REFERENCES IN FILE CA (1907 TO DATE)
58 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA
2333 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L6 ANSWER 8 OF 9 REGISTRY COPYRIGHT 2005 ACS on STN
RN 517-21-5 REGISTRY
CN 1,2-Ethanedisulfonic acid, 1,2-dihydroxy-, disodium salt (7CI, 8CI, 9CI) (CA INDEX NAME)
OTHER NAMES:

CN Disodium 1,2-dihydroxy-1,2-ethanedisulfonate
 CN Glyoxal disodium bisulfite
 CN Glyoxal-sodium bisulfite adduct (1:2)
 CN NSC 18262
 CN Sodium glyoxal bisulfite
 DR 98769-15-4, 136369-95-4, 69098-87-9, 70975-10-9, 152159-10-9, 41266-48-2,
 335081-70-4
 MF C2 H6 O8 S2 . 2 Na
 CI COM
 LC STN Files: BEILSTEIN*, BIOSIS, CA, CAOLD, CAPLUS, CASREACT, CHEMCATS,
 CHEMLIST, CSChem, IFICDB, IFIPAT, IFIUDB, MRCK*, TOXCENTER, USPAT2,
 USPATFULL
 (*File contains numerically searchable property data)
 Other Sources: EINECS**, NDSL**, TSCA**
 (**Enter CHEMLIST File for up-to-date regulatory information)
 DT.CA Caplus document type: Conference; Journal; Patent; Report
 RL.P Roles from patents: BIOL (Biological study); OCCU (Occurrence); PREP
 (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or
 reagent); USES (Uses); NORL (No role in record)
 RLD.P Roles for non-specific derivatives from patents: RACT (Reactant or
 reagent); USES (Uses)
 RL.NP Roles from non-patents: BIOL (Biological study); PREP (Preparation);
 PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES
 (Uses); NORL (No role in record)
 CRN (18381-20-9)



● 2 Na

121 REFERENCES IN FILE CA (1907 TO DATE)
 2 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA
 121 REFERENCES IN FILE CAPLUS (1907 TO DATE)
 3 REFERENCES IN FILE CAOLD (PRIOR TO 1967)


L6 ANSWER 9 OF 9 REGISTRY COPYRIGHT 2005 ACS on STN
 RN 144-62-7 REGISTRY
 CN Ethanedioic acid (9CI) (CA INDEX NAME)
 OTHER CA INDEX NAMES:
 CN Oxalic acid (8CI)
 OTHER NAMES:
 CN Aktisal
 CN Aquisal
 CN NSC 132055
 CN NSC 151956
 CN NSC 62774
 CN NSC 76990
 CN Oxagel
 CN Ultraplast Activate S 52
 FS 3D CONCORD
 DR 63504-28-9, 97993-78-7, 216451-38-6
 MF C2 H2 O4
 CI COM
 LC STN Files: ADISNEWS, AGRICOLA, ANABSTR, AQUIRE, BEILSTEIN*, BIOBUSINESS,
 BIOSIS, BIOTECHNO, CA, CABA, CANCERLIT, CAOLD, CAPLUS, CASREACT, CBNB,
 CEN, CHEMCATS, CHEMINFORMRX, CHEMLIST, CHEMSAFE, CIN, CSChem, CSNB,
 DDFU, DETHERM*, DIOGENES, DIPPR*, DRUGU, EMBASE, ENCOMPLIT, ENCOMPLIT2,
 ENCOMPPAT, ENCOMPPAT2, GMELIN*, HODOC*, HSDB*, IFICDB, IFIPAT, IFIUDB,
 IPA, MEDLINE, MRCK*, MSDS-OHS, NAPRALERT, NIOSHTIC, PDLCOM*, PIRA,
 PROMT, PS, RTECS*, SPECINFO, SYNTHLINE, TOXCENTER, TULSA, ULIDAT,
 USPAT2, USPATFULL, VETU, VTB
 (*File contains numerically searchable property data)
 Other Sources: DSL**, EINECS**, TSCA**
 (**Enter CHEMLIST File for up-to-date regulatory information)
 DT.CA Caplus document type: Book; Conference; Dissertation; Journal; Patent;
 Preprint; Report
 RL.P Roles from patents: ANST (Analytical study); BIOL (Biological study);
 FORM (Formation, nonpreparative); MSC (Miscellaneous); OCCU
 (Occurrence); PREP (Preparation); PROC (Process); PRP (Properties); RACT
 (Reactant or reagent); USES (Uses); NORL (No role in record)

Search done by Noble Jarrell

RLD.P Roles for non-specific derivatives from patents: ANST (Analytical study); BIOL (Biological study); FORM (Formation, nonpreparative); MSC (Miscellaneous); OCCU (Occurrence); PREP (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses)

RL.NP Roles from non-patents: ANST (Analytical study); BIOL (Biological study); CMBI (Combinatorial study); FORM (Formation, nonpreparative); MSC (Miscellaneous); OCCU (Occurrence); PREP (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses); NORL (No role in record)

RLD.NP Roles for non-specific derivatives from non-patents: ANST (Analytical study); BIOL (Biological study); FORM (Formation, nonpreparative); MSC (Miscellaneous); OCCU (Occurrence); PREP (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses)



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

28625 REFERENCES IN FILE CA (1907 TO DATE)
 2166 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA
 28668 REFERENCES IN FILE CAPLUS (1907 TO DATE)
 6 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

=> b wpix

FILE 'WPIX' ENTERED AT 12:51:57 ON 14 JAN 2005
 COPYRIGHT (C) 2005 THE THOMSON CORPORATION

FILE LAST UPDATED: 12 JAN 2005 <20050112/UP>
 MOST RECENT DERWENT UPDATE: 200503 <200503/DW>
 DERWENT WORLD PATENTS INDEX SUBSCRIBER FILE, COVERS 1963 TO DATE

>>> FOR A COPY OF THE DERWENT WORLD PATENTS INDEX STN USER GUIDE,
 PLEASE VISIT:
http://www.stn-international.de/training_center/patents/stn_guide.pdf <<<

>>> FOR DETAILS OF THE PATENTS COVERED IN CURRENT UPDATES, SEE
<http://thomsonderwent.com/coverage/latestupdates/> <<<

>>> FOR INFORMATION ON ALL DERWENT WORLD PATENTS INDEX USER
 GUIDES, PLEASE VISIT:
<http://thomsonderwent.com/support/userguides/> <<<

>>> NEW! FAST-ALERTING ACCESS TO NEWLY-PUBLISHED PATENT
 DOCUMENTATION NOW AVAILABLE IN DERWENT WORLD PATENTS INDEX
 FIRST VIEW - FILE WPIFV.
 FOR FURTHER DETAILS: <http://www.thomsonderwent.com/dwpifv> <<<

>>> NEW DISPLAY FORMAT HITSTR ADDED ALLOWING DISPLAY OF
 HIT STRUCTURES WITHIN THE BIBLIOGRAPHIC DOCUMENT <<<

>>> SMILES and ISOSMILES strings are no longer available as
 Derwent Chemistry Resource display fields <<<

>>> THE CPI AND EPI MANUAL CODES HAVE BEEN REVISED FROM UPDATE 200501.
 PLEASE CHECK:
<http://thomsonderwent.com/support/dwpiref/reftools/classification/code-revision/>
 FOR DETAILS. <<<

=> d all 110 tot

L10 ANSWER 1 OF 1 WPIX COPYRIGHT 2005 THE THOMSON CORP on STN
 AN 1999-405477 [34] WPIX
 DNC C1999-119773
 TI Bibenzimidazole derivatives useful for treating solid mammalian tumors or
 hematological malignancies.
 DC B02
 IN KIM, J S; LAVOIE, E J; LIU, L F; LA VOIE, E J
 PA (KIMS-I) KIM S; (LAVO-I) LAVOIE E J; (LIUL-I) LIU L F; (RUTF) UNIV RUTGERS
 STATE NEW JERSEY; (KIMJ-I) KIM J S

Search done by Noble Jarrell

CYC 85
 PI WO 9933824 A1 19990708 (199934)* EN 27 C07D403-04
 RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW NL
 OA PT SD SE SZ UG ZW
 W: AL AM AT AU AZ BA BB BG BR BY CA CH CN CU CZ DE DK EE ES FI GB GD
 GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS LT LU LV
 MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL TJ TM TR TT
 UA UG US UZ VN YU ZW
 AU 9920220 A 19990719 (199951) C07D403-04
 EP 1044199 A1 20001018 (200053) EN C07D403-04
 R: AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE
 CZ 2000002497 A3 20001213 (200103) C07D403-04
 CN 1299358 A 20010613 (200158) C07D403-04
 MX 2000006499 A1 20010201 (200168) A61K031-415
 JP 2002509858 W 20020402 (200225) 31 C07D235-06
 US 2002151575 A1 20021017 (200270) A61K031-4184 <--
 AU 753268 B 20021010 (200279) C07D403-04
 US 6667302 B2 20031223 (200408) C07D403-10 <--
 US 2004082637 A1 20040429 (200429) A61K031-4184 <--
 ADT WO 9933824 A1 WO 1998-US27822 19981230; AU 9920220 A AU
 1999-20220 19981230; EP 1044199 A1 EP 1998-965021 19981230, WO
 1998-US27822 19981230; CZ 2000002497 A3 WO 1998-US27822
 19981230, CZ 2000-2497 19981230; CN 1299358 A CN 1998-813818
 19981230; MX 2000006499 A1 MX 2000-6499 20000629; JP 2002509858 W WO
 1998-US27822 19981230, JP 2000-526506 19981230; US 2002151575 A1
 Provisional US 1997-70287P 19971231, Cont of WO 1998-US27822
 19981230, US 2001-869141 20010613; AU 753268 B AU 1999-20220
 19981230; US 6667302 B2 Provisional US 1997-70287P 19971231,
 Cont of WO 1998-US27822 19981230, US 2001-869141 20010613; US
 2004082637 A1 Provisional US 1997-70287P 19971231, Cont of
 WO 1998-US27822 19981230, Div ex US 2001-869141 20010613, US
 2003-690800 20031021
 FDT AU 9920220 A Based on WO 9933824; EP 1044199 A1 Based on WO 9933824; CZ
 2000002497 A3 Based on WO 9933824; JP 2002509858 W Based on WO 9933824; AU
 753268 B Previous Publ. AU 9920220, Based on WO 9933824; US 2004082637 A1
 Div ex US 6667302
 PRAI US 1997-70287P 19971231; US 2001-869141
 20010613; US 2003-690800 20031021
 IC ICM A61K031-415; A61K031-4184; C07D235-06; C07D403-04; C07D403-10
 ICS A61P035-00; C07D235-04; C07D403-02
 AB WO 9933824 A UPAB: 19990825
 NOVELTY - Novel bibenzimidazole derivatives are topoisomerase I inhibitors
 and effective cytotoxic agents against cancer cells, including
 drug-resistant cancer cells.
 DETAILED DESCRIPTION - Bibenzimidazole derivatives of formula (I) and
 their salts are new.
 R1, R2 = H, 1-6C alkyl, 3-6C cycloalkyl, 1-6C alkoxy, NO2, OH, 1-6C
 haloalkyl, OCF3, halo, 3-6C cycloalkyl-1-6C alkyl, 1-6C alkanoyl,
 hydroxy-1-6C alkyl, 1-6C alkoxy-carbonyl, 1-6C alkylthio, 2-6C alkanoyloxy
 or aryl or heteroaryl (both optionally substituted by 1-3 Q); or R1 + R2 =
 methylene dioxy or benzo (optionally substituted by 1-3 Q);
 Q = 1-6C alkyl, 3-6C cycloalkyl, 1-6C alkoxy, NO2, OH, 1-6C halo
 alkyl, OCF3, 3-6C cycloalkyl-1-6C alkyl, 1-6C alkanoyl, hydroxy-1-6C
 alkyl, 1-6C alkoxy-carbonyl, 1-6C alkylthio, 2-6C alkanoyloxy or halo;
 R3 = H, 1-6C alkyl, 3-6C cycloalkyl, 1-6C alkoxy, NO2, OH, halo- 1-6C
 alkyl, OCF3, 3-6C cycloalkyl-1-6C alkyl, 1-6C alkanoyl, hydroxy-1-6C
 alkyl, 1-6C alkoxy-carbonyl, 1-6C alkylthio, 2-6C alkanoyl-oxy or halo;
 R4 + R5 = 3-5 membered saturated or unsaturated chain comprising
 non-peroxide oxygen, sulfur, N(X) and carbon, optionally substituted by
 oxo;
 X = absent or is H, O, 1-4C alkyl, phenyl or benzyl; in which at
 least one (e.g. 1 or 2) of the chain members is NH; provided that R4 + R5
 are not NH-CH=N.
 ACTIVITY - Cytostatic. 5-Phenyl-2'-(benzotriazol-5-yl)-
 bibenzimidazole showed in vitro cytotoxicity against RPMI 8402 cancer
 cells and camptothecin resistant CPT-K5 cells with IC50 values of 0.47 and
 0.47 microns, respectively.
 MECHANISM OF ACTION - Topoisomerase-I Inhibitor.
 USE - (I) are potent topoisomerase I poisons. They exhibit cytotoxic
 activity against RPMI 8402 cancer cells and camptothecin resistant CPT-K5
 cells. (I) are useful as cytotoxic agents for the treatment of cancers,
 and in particular, solid mammalian tumors or hematological malignancies.
 (I) are also useful as pharmacological tools for in vitro and in vivo
 study of topoisomerase function and activity.
 Dwg.0/2
 FS CPI

FA AB; GI; DCN
MC CPI: B06-D05; B06-H; B12-K04; B14-D09; B14-H01

=> b home
FILE 'HOME' ENTERED AT 12:52:06 ON 14 JAN 2005

=>

```
=> b reg
FILE 'REGISTRY' ENTERED AT 13:33:47 ON 14 JAN 2005
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2005 American Chemical Society (ACS)
```

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

```
STRUCTURE FILE UPDATES: 12 JAN 2005 HIGHEST RN 812631-13-3
DICTIONARY FILE UPDATES: 12 JAN 2005 HIGHEST RN 812631-13-3
```


TSCA INFORMATION NOW CURRENT THROUGH MAY 21, 2004

Please note that search-term pricing does apply when conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more information enter HELP PROP at an arrow prompt in the file or refer to the file summary sheet on the web at:
<http://www.cas.org/ONLINE/DBSS/registryss.html>

```
=> d que stat l15
L13 STR
```



```
VAR G1=20/21/24/25
VAR G2=C/N/O/S
REP G3=(1-3) A
NODE ATTRIBUTES:
DEFAULT MLEVEL IS ATOM
DEFAULT ECLEVEL IS LIMITED
```

```
GRAPH ATTRIBUTES:
RSPEC 10
NUMBER OF NODES IS 28
```

```
STEREO ATTRIBUTES: NONE
L15 83 SEA FILE=REGISTRY SSS FUL L13
```

```
100.0% PROCESSED 1243 ITERATIONS 83 ANSWERS
SEARCH TIME: 00.00.01
```

```
=> d his
```

(FILE 'HOME' ENTERED AT 12:47:52 ON 14 JAN 2005)

FILE 'HCAPLUS' ENTERED AT 12:47:58 ON 14 JAN 2005

```
L1 1 (US20040082637 OR US6667302 OR US20020151575)/PN
E W01998-US27822/AP,PRN
L2 1 W01998-US27822/AP,PRN
E US1997-070287/AP,PRN
L3 1 US1997-070287/AP,PRN
L4 1 L1-3
```

FILE 'REGISTRY' ENTERED AT 12:50:28 ON 14 JAN 2005

FILE 'HCAPLUS' ENTERED AT 12:50:31 ON 14 JAN 2005

```
L5 TRA L4 1- RN : 9 TERMS
```

Search done by Noble Jarrell

L6 FILE 'REGISTRY' ENTERED AT 12:50:31 ON 14 JAN 2005
9 SEA L5

L7 FILE 'WPIX' ENTERED AT 12:50:34 ON 14 JAN 2005
L8 1 (US20040082637 OR US6667302 OR US20020151575)/PN
L9 1 WO1998-US27822/AP,PRN
L10 1 US1997-070287P/AP,PRN
L10 1 L7-9

L11 FILE 'REGISTRY' ENTERED AT 12:54:21 ON 14 JAN 2005
L12 STR
L13 STR L11
L14 STR L12
L15 4 L13
83 L13 FULL
SAV TEM MIT800F0/A L15

L16 FILE 'HCAPLUS' ENTERED AT 13:25:23 ON 14 JAN 2005
E LAVOIE E/AU
195 E3-4,E6-7
E KIM J/AU
L17 1629 E3,E42
E KIM JUNG/AU
L18 16 E3
E KIM JUNG SUN/AU
L19 55 E3
E LIU L/AU
L20 635 E3,E10
E LIU LEROY/AU
L21 206 E3-5
L22 28252 RUTGERS/CS,PA
L23 QUE (?CANCER? OR ?TUMOR? OR ?TUMOUR? OR ?CARCINOGEN? OR ?MALIGN
L24 36 L15
L25 16 L24 AND L16-22
L26 20 L24 NOT L25
L27 3 L26 AND L23
L28 17 L26 NOT L27

=> b hcap
FILE 'HCAPLUS' ENTERED AT 13:34:13 ON 14 JAN 2005
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 14 Jan 2005 VOL 142 ISS 3
FILE LAST UPDATED: 12 Jan 2005 (20050112/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d all fhistr 125 tot


L25 ANSWER 1 OF 16 HCAPLUS COPYRIGHT 2005 ACS on STN
AN 2003:941009 HCAPLUS
DN 140:280814
ED Entered STN: 03 Dec 2003
TI Effects of topoisomerases inhibitors protoberberine on Leishmania donovani growth, macrophage function, and infection
AU Marquis, Jean-Francois; Makhey, Darshan; LaVoie, Edmond J.; Olivier, Martin
CS Departement de Biologie Medicale, Faculte de Medecine, Centre de Recherche en Infectiologie du CHUQ, Universite Laval, Sainte-Foy, QC, G1V 4G2, Can.
SO Journal of Parasitology (2003), 89(5), 1048-1052
CODEN: JOPAA2; ISSN: 0022-3395

Search done by Noble Jarrell

PB American Society of Parasitologists
 DT Journal
 LA English
 CC 1-5 (Pharmacology)
 AB DNA topoisomerases play a pivotal role in the regulation of cell division. Inhibition of Leishmania spp. topoisomerases represents an alternative to control parasite growth. Cancer research led to the development of several potent topoisomerase inhibitors such as topoisomerase 1, topoisomerase 17, or both (monobenzimidazole, terbenzimidazole, and protoberberine alkaloid-related compds.) that are effective antitumor agents. In the present study, we evaluated the efficacy of these compds. against Leishmania spp. growth in vitro. Some protoberberine compds. showed pronounced antileishmanial activity and were selected for further anal. in macrophages. These compds. did not affect macrophage viability and only slightly reduced macrophage nitric oxide generation in response to interferon- γ . Moreover, exposure of infected macrophages to these compds. significantly reduced parasite loads. Collectively, our data suggest that protoberberine-related compds. have powerful antileishmanial action and that minor structural variations among them can substantially improve their activity to restrict Leishmania spp. infection in vitro.

ST topoisomerase inhibitor protoberberine deriv Leishmania donovani
 IT Protozoacides
 (leishmanicides; topoisomerase inhibitors protoberberine derivs. effects on Leishmania donovani growth, macrophage function, and infection)
 IT Leishmania donovani
 Macrophage
 Phagocyte
 (topoisomerase inhibitors protoberberine derivs. effects on Leishmania donovani growth, macrophage function, and infection)
 IT 10102-43-9, Nitric oxide, biological studies
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (topoisomerase inhibitors protoberberine derivs. effects on Leishmania donovani growth, macrophage function, and infection)
 IT 483-15-8 2086-83-1 3486-67-7 6872-81-7 17388-19-1 19716-66-6
 35989-93-6 96954-35-7 167959-21-9, Terbenzimidazole
 180077-22-9 180077-24-1 180077-31-0 180077-32-1 180077-33-2
 286000-57-5 675881-79-5 675881-80-8 675881-81-9 675881-82-0
 675881-83-1
 RL: DMA (Drug mechanism of action); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (topoisomerase inhibitors protoberberine derivs. effects on Leishmania donovani growth, macrophage function, and infection)
 IT 19716-69-9D, Protoberberine, derivs.
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (topoisomerase inhibitors protoberberine derivs. effects on Leishmania donovani growth, macrophage function, and infection)
 RE.CNT 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS RECORD
 RE
 (1) Broccoli, S; Nucleic Acids Research 1999, V27, P2745 HCAPLUS
 (2) Cheesman, S; Parasitology Today 2000, V16, P277 HCAPLUS
 (3) Chen, A; Cancer Research 1993, V53, P1332 HCAPLUS
 (4) Chen, A; Proceedings of the National Academy of Sciences of the United States of America 1993, V90, P8131 HCAPLUS
 (5) Das, A; Nucleic Acids Research 2001, V29, P1844 HCAPLUS
 (6) Evans, T; Journal of Immunology 1993, V151, P907 HCAPLUS
 (7) Herwaldt, B; Lancet 1999, V354, P1191 MEDLINE
 (8) Kim, J; Bioorganic and Medicinal Chemistry 1996, V4, P621 HCAPLUS
 (9) Kim, J; Journal of Medicinal Chemistry 1997, V40, P2818 HCAPLUS
 (10) Liu, L; Annual Review of Biochemistry 1989, V58, P351 HCAPLUS
 (11) Makhey, D; Bioorganic and Medicinal Chemistry 1996, V4, P781 HCAPLUS
 (12) Makhey, D; Medicinal Chemistry Research 1995, V5, P1 HCAPLUS
 (13) Mansfield, J; Immunology of infectious diseases 2002, P379 HCAPLUS
 (14) Marquis, J; Parasitology 2003, V126, P21 HCAPLUS
 (15) Matte, C; European Journal of Immunology 2000, V30, P2555 HCAPLUS
 (16) Olivier, M; Infection and Immunity 1987, V55, P467 MEDLINE
 (17) Olivier, M; Journal of Biological Chemistry 1998, V273, P13944 HCAPLUS
 (18) Radzioch, D; Journal of Leukocyte Biology 1991, V50, P263 HCAPLUS
 (19) Rangarajan, M; Bioorganic and Medicinal Chemistry 2000, V8, P1371 HCAPLUS
 (20) Sun, Q; Bioorganic and Medicinal Chemistry Letters 1994, V4, P2871 HCAPLUS
 (21) Sun, Q; Journal of Medicinal Chemistry 1995, V38, P3638 HCAPLUS
 (22) Villa, H; Journal of Biological Chemistry 2003, V6, P3521
 (23) Wang, J; Annual Review of Biochemistry 1985, V54, P665 MEDLINE
 (24) Wang, J; Annual Review of Biochemistry 1996, V65, P635 HCAPLUS

(25) Wang, L; Chemical Research in Toxicology 1996, V9, P75 HCAPLUS
 (26) White, T; Journal of Biological Chemistry 1988, V263, P16977 HCAPLUS
 IT 167959-21-9, Terbenzimidazole
 RL: DMA (Drug mechanism of action); PAC (Pharmacological activity); THU
 (Therapeutic use); BIOL (Biological study); USES (Uses)
 (topoisomerase inhibitors protoberberine derivs. effects on Leishmania
 donovani growth, macrophage function, and infection)
 RN 167959-21-9 HCAPLUS
 CN 2,5':2',5''-Ter-1H-benzimidazole (9CI) (CA INDEX NAME)




L25 ANSWER 2 OF 16 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 2003:679625 HCAPLUS
 DN 140:280441
 ED Entered STN: 31 Aug 2003
 TI Defining the molecular interactions that are important for the poisoning
 of human topoisomerase I by benzimidazoles and terbenzimidazoles
 AU Pilch, Daniel S.; Liu, Hsing-Yin; Li, Tsai-Kun; Kerrigan, John E.;
 LaVoie, Edmond J.; Barbieri, Christopher M.
 CS Germany
 SO Small Molecule DNA and RNA Binders (2003), Volume 2, 576-608. Editor(s):
 Demeunynck, Martine; Bailly, Christian; Wilson, W. David. Publisher:
 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, Germany.
 CODEN: 69EKGP; ISBN: 3-527-30595-5
 DT Conference; General Review
 LA English
 CC 1-0 (Pharmacology)
 AB A review focuses on the mol. interactions that are important in the
 poisoning of human topoisomerase type I by antitumor benzimidazole-containing
 compds., with emphasis on derivs. that contain either one or three
 benzimidazole functionalities.
 ST review human topoisomerase I poisoning benzimidazole terbenzimidazole;
 antitumor benzimidazole topoisomerase I poisoning review
 IT Antitumor agents
 Human
 Molecular association
 Neoplasm
 (defining the mol. interactions that are important for poisoning of
 human topoisomerase I by antitumor benzimidazoles and
 terbenzimidazoles)
 IT 143180-75-0, DNA Topoisomerase I
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (defining the mol. interactions that are important for poisoning of
 human topoisomerase I by antitumor benzimidazoles and
 terbenzimidazoles)
 IT 51-17-2D, Benzimidazole, analogs 167959-21-9D, Terbenzimidazole,
 analogs
 RL: DMA (Drug mechanism of action); PAC (Pharmacological activity); THU
 (Therapeutic use); BIOL (Biological study); USES (Uses)
 (defining the mol. interactions that are important for poisoning of
 human topoisomerase I by antitumor benzimidazoles and
 terbenzimidazoles)
 RE.CNT 122 THERE ARE 122 CITED REFERENCES AVAILABLE FOR THIS RECORD
 RE
 (1) Baguley, B; Nucleic Acids Res 1978, V5, P161 HCAPLUS
 (2) Bailly, C; Biochemistry 1997, V36, P3917 HCAPLUS
 (3) Bauer, W; J Mol Biol 1968, V33, P141 HCAPLUS
 (4) Baxter, C; Proteins 1998, V33, P367 HCAPLUS
 (5) Been, M; Nucleic Acids Res 1984, V12, P3097 HCAPLUS
 (6) Bharti, A; J Biol Chem 1996, V271, P1993 HCAPLUS
 (7) Burke, T; Anal Biochem 1993, V212, P285 HCAPLUS
 (8) Cantor, C; Biophysical Chemistry, Part I: The Conformation of Biological
 Macromolecules 1980
 (9) Caserta, M; Nucleic Acids Res 1989, V17, P8463 HCAPLUS
 (10) Castano, I; Genes Dev 1996, V10, P2564 HCAPLUS
 (11) Chen, A; Cancer Res 1993, V53, P1332 HCAPLUS

- (12) Chen, A; Proc Natl Acad Sci USA 1993, V90, P8131 HCAPLUS
- (13) Chen, H; J Mol Biol 1996, V258, P457 HCAPLUS
- (14) Chen, J; Nucleic Acids Res 1988, V16, P6803 HCAPLUS
- (15) Chen, Y; Annu Rev Pharmacol Toxicol 1994, V34, P191
- (16) Chourpa, I; Biochim Biophys Acta 1998, V1379, P353 HCAPLUS
- (17) Christiansen, K; J Biol Chem 1993, V268, P9690 HCAPLUS
- (18) Clark, G; Biochemistry 1996, V35, P13745 HCAPLUS
- (19) Crawford, L; J Mol Biol 1967, V25, P23 HCAPLUS
- (20) Cummings, J; Biochem Pharmacol 2002, V63, P607 HCAPLUS
- (21) D'Arpa, P; Proc Natl Acad Sci USA 1988, V85, P2543 HCAPLUS
- (22) Edwards, T; J Biol Chem 2000, V275, P36181 HCAPLUS
- (23) Fan, Y; J Med Chem 1998, V41, P2216 HCAPLUS
- (24) Fassberg, J; J Pharm Sci 1992, V81, P676 HCAPLUS
- (25) Fede, A; Biochemistry 1991, V30, P11377 HCAPLUS
- (26) Fede, A; Structure 1993, V1, P177 HCAPLUS
- (27) Gatto, B; Cancer Res 1996, V56, P2795 HCAPLUS
- (28) Geacintov, N; Biochemistry 1978, V17, P5256 HCAPLUS
- (29) Geacintov, N; Biochemistry 1987, V26, P3087 HCAPLUS
- (30) Giovanella, B; Cancer Res 1991, V51, P3052 HCAPLUS
- (31) Giovanella, B; Science 1989, V246, P1046 HCAPLUS
- (32) Hagerman, P; Biochemistry 1985, V24, P7033 HCAPLUS
- (33) Hagerman, P; Nature 1986, V321, P449 HCAPLUS
- (34) Haluska, P; Advan Enzyme Regul 1998, V38, P253 HCAPLUS
- (35) Haluska, P; Nucleic Acids Res 1998, V26, P1841 HCAPLUS
- (36) Haluska, P; Nucleic Acids Res 1999, V27, P2538 HCAPLUS
- (37) Holden, J; Arch Biochem Biophys 1999, V370, P66 HCAPLUS
- (38) Hsiang, Y; J Biol Chem 1985, V260, P14873 HCAPLUS
- (39) Husain, I; Proc Natl Acad Sci USA 1988, V85, P2558 HCAPLUS
- (40) Jin, S; Bioorg Med Chem Lett 2000, V10, P719 HCAPLUS
- (41) Jones, R; Biochemistry 1986, V25, P7431 HCAPLUS
- (42) Juan, C; Proc Natl Acad Sci USA 1988, V85, P8910 HCAPLUS
- (43) Kanzawa, F; Cancer Res 1995, V55, P2806 HCAPLUS
- (44) Keller, W; Proc Natl Acad Sci USA 1975, V72, P4876 HCAPLUS
- (45) Kerrigan, J; Biochemistry 2001, V40, P9792 HCAPLUS
- (46) Kim, J; Bioorg Med Chem 1996, V4, P621 HCAPLUS
- (47) Kim, J; Bioorg Med Chem 1998, V6, P163 HCAPLUS
- (48) Kim, J; J Med Chem 1997, V40, P2818 HCAPLUS
- (49) Knab, A; J Biol Chem 1993, V268, P22322 HCAPLUS
- (50) Koo, H; Biochemistry 1990, V29, P4227 HCAPLUS
- (51) Koo, H; Nature 1986, V320, P501 HCAPLUS
- (52) Krogh, S; Nucleic Acids Res 1991, V19, P1235 HCAPLUS
- (53) Lanza, A; J Biol Chem 1996, V271, P6978 HCAPLUS
- (54) Lee, M; Proc Natl Acad Sci USA 1993, V90, P6656 HCAPLUS
- (55) Levene, S; Biochemistry 1986, V25, P3988 HCAPLUS
- (56) Li, T; Annu Rev Pharmacol Toxicol 2001, V41, P53 HCAPLUS
- (57) Li, T; Biochemistry 2000, V39, P7107 HCAPLUS
- (58) Liu, L; Annu Rev Biochem 1989, V58, P351 HCAPLUS
- (59) Liu, L; Biochim Biophys Acta 1975, V395, P405 HCAPLUS
- (60) Makhey, D; Med Chem Res 1995, V5, P1 HCAPLUS
- (61) Marini, J; J Biol Chem 1984, V259, P8974 HCAPLUS
- (62) Marini, J; Proc Natl Acad Sci USA 1982, V79, P7664 HCAPLUS
- (63) McGhee, J; J Mol Biol 1974, V86, P469 HCAPLUS
- (64) Mi, Z; Biochemistry 1994, V33, P12540 HCAPLUS
- (65) Mills, J; Biochemistry 1994, V33, P1797 HCAPLUS
- (66) Moon, J; Biopolymers 1996, V38, P593 HCAPLUS
- (67) Morham, S; Mol Cell Biol 1996, V16, P6804 HCAPLUS
- (68) Munoz, M; Spectrochim Acta Part A 2001, V57, P1049
- (69) Norden, B; Appl Spectrosc Rev 1978, V14, P157 HCAPLUS
- (70) Norden, B; Q Rev Biophys 1992, V25, P51 HCAPLUS
- (71) Pantazis, P; Cancer Res 1992, V52, P3980 HCAPLUS
- (72) Pantazis, P; Cancer Res 1993, V53, P1577 HCAPLUS
- (73) Pantazis, P; Int J Cancer 1993, V53, P863 HCAPLUS
- (74) Park, Y; Proc Natl Acad Sci USA 1991, V88, P1551 HCAPLUS
- (75) Parkinson, G; Nature Struct Biol 1996, V3, P837 HCAPLUS
- (76) Patel, S; Investig New Drugs 1991, V9, P53 MEDLINE
- (77) Pearl, D; Science 1985, V227, P1304
- (78) Perez-Stable, C; Nucleic Acids Res 1988, V16, P7975 HCAPLUS
- (79) Pilch, D; Biochemistry 1995, V34, P9962 HCAPLUS
- (80) Pilch, D; Biochemistry 1997, V36, P12542 HCAPLUS
- (81) Pilch, D; Drug Des Disc 1996, V13(3-4), P115 HCAPLUS
- (82) Pilch, D; Proc Natl Acad Sci USA 1997, V94, P13565 HCAPLUS
- (83) Pjura, P; J Mol Biol 1987, V197, P257 HCAPLUS
- (84) Pommier, Y; Biochim Biophys Acta 1998, V1400, P83 HCAPLUS
- (85) Pommier, Y; Drug Resis Upd 1999, V2, P307 HCAPLUS
- (86) Porter, S; Nucleic Acids Res 1989, V17, P8521 HCAPLUS
- (87) Qiu, D; J Phys Chem A 1997, V101, P3005 HCAPLUS

- (88) Rangarajan, M; Bioorg Med Chem 2000, V8, P1371 HCAPLUS
 (89) Redinbo, M; Science 1998, V279, P1504 HCAPLUS
 (90) Revet, B; Nature New Biol 1971, V229, P10 HCAPLUS
 (91) Robinson, H; Gene 1994, V149, P179 HCAPLUS
 (92) Saleem, A; Ann NY Acad Sci 2000, V922, P46 HCAPLUS
 (93) Sanders, M; Biochem Pharmacol 1997, V56, P1157
 (94) Shaiu, W; Mol Cell Biol 1998, V18, P4358 HCAPLUS
 (95) Shen, C; J Mol Biol 1990, V212, P67 HCAPLUS
 (96) Sim, S; Biochemistry 1997, V36, P13285 HCAPLUS
 (97) Sim, S; Biochemistry 2000, V39, P9928 HCAPLUS
 (98) Simmons, D; Virology 1996, V222, P365 HCAPLUS
 (99) Sobell, H; J Mol Biol 1972, V68, P21 HCAPLUS
 (100) Spink, N; Nucleic Acids Res 1994, V22, P1607 HCAPLUS
 (101) Stevnsner, T; J Biol Chem 1989, V264, P10110 HCAPLUS
 (102) Stewart, L; The Journal of Biological Chemistry 1996, V271, P7602 HCAPLUS
 (103) Strauss, J; Science 1994, V266, P1829 HCAPLUS
 (104) Sun, Q; J Med Chem 1995, V38, P3638 HCAPLUS
 (105) Svejstrup, J; J Mol Biol 1991, V222, P669 HCAPLUS
 (106) Takahashi, T; Jpn J Cancer Res 1997, V88, P1211 HCAPLUS
 (107) Takusagawa, F; Cold Spring Harbor Symp Quant Biol 1983, V47, P315
 (108) Teng, M; Nucleic Acids Res 1988, V16, P2671 HCAPLUS
 (109) Vega, M; Eur J Biochem 1994, V222, P721 HCAPLUS
 (110) Wall, M; J Am Chem Soc 1966, V88, P3888 HCAPLUS
 (111) Wang, J; Annu Rev Biochem 1996, V65, P635 HCAPLUS
 (112) Wang, J; J Mol Biol 1974, V89, P783 HCAPLUS
 (113) Wang, L; Chem Res Toxicol 1993, V6, P813 HCAPLUS
 (114) Wani, M; J Med Chem 1986, V29, P2358 HCAPLUS
 (115) Wells, R; J Mol Biol 1970, V49, P319 HCAPLUS
 (116) Wilson, R; Biopolymers 1978, V17, P1235 HCAPLUS
 (117) Wilson, W; Adv Pharmacol Chemother 1981, V18, P177 HCAPLUS
 (118) Wu, H; Nature 1984, V308, P509 HCAPLUS
 (119) Xu, Z; Biochemistry 1998, V37, P3558 HCAPLUS
 (120) Yamashita, Y; Biochemistry 1992, V31, P12069 HCAPLUS
 (121) Yoshinari, T; Cancer Res 1993, V53, P490 HCAPLUS
 (122) Zhang, C; Dev Biol 2000, V222, P27 HCAPLUS

IT 167959-21-9D, Terbenzimidazole, analogs
 RL: DMA (Drug mechanism of action); PAC (Pharmacological activity); THU
 (Therapeutic use); BIOL (Biological study); USES (Uses)
 (defining the mol. interactions that are important for poisoning of
 human topoisomerase I by antitumor benzimidazoles and
 terbenzimidazoles)

RN 167959-21-9 HCAPLUS
 CN 2,5':2',5''-Ter-1H-benzimidazole (9CI) (CA INDEX NAME)



L25 ANSWER 3 OF 16 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 2000:840668 HCAPLUS
 DN 134:95134
 ED Entered STN: 01 Dec 2000
 TI Topoisomerase I inhibition and cytotoxicity of 5-bromo- and
 5-phenylterbenzimidazoles
 AU Rangarajan, Meera; Kim, Jung Sun; Sim, Sai-Peng; Liu, Angela;
 Liu, Leroy F.; LaVoie, Edmond J.
 CS Department of Pharmaceutical Chemistry, Rutgers, The State
 University of New Jersey, Piscataway, NJ, 08854, USA
 SO Bioorganic & Medicinal Chemistry (2000), 8(11), 2591-2600
 CODEN: BMECEP; ISSN: 0968-0896
 PB Elsevier Science Ltd.
 DT Journal
 LA English
 CC 1-3 (Pharmacology)
 Section cross-reference(s): 7, 28
 OS CASREACT 134:95134
 AB Topoisomerase I is an enzyme that is essential for maintaining the
 three-dimensional structure of DNA during the processes of transcription,
 translation and mitosis. With the introduction of new clin. agents that

are effective in poisoning topoisomerase I, this enzyme has proved to be an attractive mol. target in the development of anticancer drugs. Several terbenzimidazoles have been identified as potent topoisomerase I poisons. Structure-activity data on various terbenzimidazoles have revealed that the presence of lipophilic substituents at the 5-position of various terbenzimidazoles correlates with enhanced cytotoxicity. While the effect of having substituents at both the 5- and 6-positions had not been evaluated, previous studies did indicate that the presence of a fused benzo-ring at the 5,6-position results in a significant decrease in topoisomerase I poisoning activity and cytotoxicity. In the present study we investigated whether substituents at both the 5- and 6-positions of varied terbenzimidazoles would allow for retention of topo I poisoning activity. The 6-bromo, 6-methoxy, or 6-Ph derivs. of both 5-bromo- and 5-phenylterbenzimidazole were synthesized and evaluated for topo I poisoning activity, as well as their cytotoxicity toward human lymphoblastoma cells. The data indicate that such derivs. do retain similar topo I poisoning activity and possess cytotoxicity equivalent to either 5-bromo- or 5-phenylterbenzimidazole. Significant enhancement in the topoisomerase I poisoning activity and cytotoxicity of 5-phenylterbenzimidazole is observed when the 2"-position is substituted with either a chloro or trifluoromethyl substituent. The influence of such substituents on the biol. activity of 5,6-dibromoterbenzimidazole (I) was also explored. In the case of either 2"-chloro-5,6-dibromoterbenzimidazole or 2"-trifluoromethyl-5,6-dibromoterbenzimidazole (II), topoisomerase I poisoning was not enhanced relative to I. While cytotoxicity toward RPMI 8402 was also not significantly affected comparative studies performed against several solid human tumor cell lines did reveal a significant increase in cytotoxicity observed for II as compared to I.

ST terbenzimidazole prepn topoisomerase inhibition cytotoxicity structure

IT Antitumor agents

Structure-activity relationship

(synthesis, topoisomerase I inhibition and cytotoxicity of terbenzimidazoles)

IT 167959-27-5P 185199-36-4P 237429-45-7P

237429-52-6P 237429-53-7P 237429-54-8P

237429-55-9P 237429-56-0P 237429-57-1P

237429-58-2P 237429-59-3P 288579-81-7P

319916-61-5P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(synthesis, topoisomerase I inhibition and cytotoxicity of terbenzimidazoles)

IT 143180-75-0

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(synthesis, topoisomerase I inhibition and cytotoxicity of terbenzimidazoles)

IT 875-51-4, 4-Bromo-2-nitroaniline 1679-18-1, p-Chlorophenylboronic acid

6943-69-7 17626-40-3 35998-98-2 75293-97-9, 3,4-Dibromo-6-

nitroaniline 106429-45-2 106429-59-8 167959-20-8 288579-82-8

RL: RCT (Reactant); RACT (Reactant or reagent)

(synthesis, topoisomerase I inhibition and cytotoxicity of terbenzimidazoles)

IT 1575-37-7P 31433-98-4P 49764-63-8P 108447-01-4P 117878-22-5P,

[1,1':2',1''-Terphenyl]-4',5'-diamine 185199-45-5P 237429-70-8P

237429-71-9P 237429-73-1P 237429-74-2P 237429-75-3P 237429-76-4P

237429-77-5P 237429-78-6P 319916-62-6P 319916-63-7P 319916-64-8P

319916-65-9P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(synthesis, topoisomerase I inhibition and cytotoxicity of terbenzimidazoles)


RE.CNT 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Abdullah, R; US 4264600 1981 HCAPLUS
- (2) Beerman, T; Biochim Biophys Acta 1992, V1131, P53 HCAPLUS
- (3) Carmichael, J; Cancer Res 1987, V47, P936 HCAPLUS
- (4) Chen, A; Annu Rev Pharmacol Toxicol 1994, V34, P191 HCAPLUS
- (5) Chen, A; Cancer Res 1993, V53, P1332 HCAPLUS
- (6) Chen, A; Proc Natl Acad Sci USA 1993, V90, P8131 HCAPLUS
- (7) Clark, R; J Am Chem Soc 1958, V80, P1657 HCAPLUS
- (8) Corey, E; Tet Lett 1979, V5, P399
- (9) Denizot, F; J Immunol Methods 1986, V89, P271 MEDLINE
- (10) Hsiang, Y; J Biol Chem 1985, V260, P14873 HCAPLUS

Search done by Noble Jarrell

- (11) Kim, J; Bioorg Med Chem 1998, V6, P163 HCAPLUS
 (12) Kim, J; J Med Chem 1997, V40, P2818 HCAPLUS
 (13) Liu, L; Annu Rev Biochem 1989, V58, P351 HCAPLUS
 (14) Liu, L; J Biol Chem 1983, V258, P15365 HCAPLUS
 (15) Miyaura, N; Chem Rev 1995, V95, P2457 HCAPLUS
 (16) Mosman, T; J Immunol Methods 1983, V65, P55
 (17) Phillips, M; J Chem Soc 1928, P2393 HCAPLUS
 (18) Pommier, Y; Biochim Biophys Acta 1998, V1400, P83 HCAPLUS
 (19) Preston, P; Benzimidazoles. Benzimidazoles and Congeneric Tricyclic Compounds 1981, Part 1, P1
 (20) Rangarajan, M; Bioorg Med Chem Lett, Submitted
 (21) Stephens, F; J Chem Soc 1950, V1722
 (22) Sun, Q; J Med Chem 1995, V38, P3638 HCAPLUS
 (23) Wang, J; Annu Rev Biochem 1996, V65, P635 HCAPLUS
 (24) Xu, Z; Biochemistry 1998, V37, P3558 HCAPLUS
 IT 167959-27-5P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (synthesis, topoisomerase I inhibition and cytotoxicity of terbenzimidazoles)
 RN 167959-27-5 HCAPLUS
 CN 2,5':2',5''-Ter-1H-benzimidazole, 5-phenyl- (9CI) (CA INDEX NAME)



- L25 ANSWER 4 OF 16 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 2000:454831 HCAPLUS
 DN 133:171757
 ED Entered STN: 06 Jul 2000
 TI 2''-Substituted 5-phenylterbenzimidazoles as topoisomerase I poisons
 AU Rangarajan, M.; Kim, J. S.; Jin, S.; Sim, S.-P.; Liu, A.; Pilch, D. S.; Liu, L. F.; LaVoie, E. J.
 CS Department of Pharmaceutical Chemistry, Rutgers, The State University of New Jersey, Piscataway, NJ, 08854, USA
 SO Bioorganic & Medicinal Chemistry (2000), 8(6), 1371-1382
 CODEN: BMECEP; ISSN: 0968-0896
 PB Elsevier Science Ltd.
 DT Journal
 LA English
 CC 1-3 (Pharmacology)
 Section cross-reference(s): 28
 AB 5-Phenylterbenzimidazole (1) is active as a topoisomerase I poison (topo I) and is cytotoxic to human tumor cells. No cross-resistance was observed for 1 when it was evaluated against the camptothecin-resistant cell line, CPT-K5. Derivs. of 1 substituted at the 2''-position, however, did exhibit cross-resistance to this cell line. The basis for the resistance of this cell line towards CPT is that it possesses a mutant form of topo I. These results suggest that substituents at the 2''-position may be in proximity to the wild-type enzyme. Therefore, we hypothesized that terbenzimidazoles with 2''-substituents could be capable of interacting with the enzyme and thereby influence activity within this class of topo I poisons. 5-Phenylterbenzimidazoles with a hydroxy, hydroxymethyl, mercapto, amino, N-benzoylaminomethyl, chloro, and trifluoromethyl group at the 2''-position were synthesized. In addition, several 2''-ethyl-5-phenylterbenzimidazoles were prepared containing either a methoxy, hydroxy, amino, or N-acetyl amino group at the 2-position of the Et side-chain. These 2''-substituted 5-phenylterbenzimidazoles were evaluated as topo I poisons and for cytotoxic activity. The presence of a strong electron-withdrawing group at the 2''-position, such as a chloro or trifluoromethyl group, did enhance both topo I poisoning activity and cytotoxicity. Studies on the relative DNA binding affinity of 1 to its 2''-amino and 2''-trifluoromethyl derivs. did exhibit a correlation with their relative differences in biol. activity.
 ST phenylterbenzimidazole deriv prepn antitumor topoisomerase inhibitor
 IT DNA
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(binding; preparation of phenylterbenzimidazoles as topoisomerase I poisons)

IT Antitumor agents
Structure-activity relationship
(preparation of phenylterbenzimidazoles as topoisomerase I poisons)

IT 237429-44-6P 237429-45-7P
RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses)
(preparation of phenylterbenzimidazoles as topoisomerase I poisons)

IT 167959-27-5
RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)
(preparation of phenylterbenzimidazoles as topoisomerase I poisons)

IT 237429-50-4P 237429-51-5P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)
(preparation of phenylterbenzimidazoles as topoisomerase I poisons)

IT 192879-73-5P 192879-74-6P 192879-75-7P
237429-42-4P 237429-43-5P 237429-46-8P
237429-47-9P 237429-48-0P 237429-49-1P
288579-81-7P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of phenylterbenzimidazoles as topoisomerase I poisons)

IT 143180-75-0
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
(preparation of phenylterbenzimidazoles as topoisomerase I poisons)


IT 57-13-6, Urea, reactions 76-05-1, Trifluoroacetic acid, reactions 79-14-1, reactions 107-95-9, .beta.-Alanine 140-89-6, Ethylxanthic acid potassium salt 495-69-2, Hippuric acid 506-68-3, Cyanogen bromide 4324-37-2, 2-Methoxypropionic acid 17626-40-3 35998-98-2, 3,4-Dinitrobenzaldehyde 192879-70-2
RL: RCT (Reactant); RACT (Reactant or reagent)
(preparation of phenylterbenzimidazoles as topoisomerase I poisons)

IT 3671-61-2P 106429-45-2P 106429-59-8P 192879-72-4P 221289-88-9P
230308-95-9P 237429-60-6P 237429-61-7P 237429-62-8P 237429-63-9P
237429-64-0P 237429-65-1P 237429-66-2P 237429-67-3P 237429-68-4P
237429-69-5P 288579-82-8P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation of phenylterbenzimidazoles as topoisomerase I poisons)

RE.CNT 37 THERE ARE 37 CITED REFERENCES AVAILABLE FOR THIS RECORD
RE

- (1) Beerman, T; Biochim Biophys Acta 1992, V1131, P53 HCAPLUS
- (2) Bjornsti, M; Cancer Res 1989, V49, P6318 HCAPLUS
- (3) Breslauer, K; J Meth Enzymol 1995, V259, P221 HCAPLUS
- (4) Carmichael, J; Cancer Res 1987, V47, P936 HCAPLUS
- (5) Chen, A; Annu Rev Pharmacol Toxicol 1994, V34, P191 HCAPLUS
- (6) Chen, A; Cancer Res 1993, V53, P1332 HCAPLUS
- (7) Chen, A; Proc Natl Acad Sci 1993, V90, P8131 HCAPLUS
- (8) Clark, G; Biochemistry 1996, V35, P13745 HCAPLUS
- (9) Crothers, D; Biopolymers 1971, V10, P2147 HCAPLUS
- (10) Denizot, F; J Immunol Meth 1986, V89, P271 MEDLINE
- (11) Hertzberg, R; Biochemistry 1989, V28, P4629 HCAPLUS
- (12) Hsiang, Y; J Biol Chem 1985, V260, P14873 HCAPLUS
- (13) Hughes, G; J Proc Roy Soc N S Wales 1938, V71, P209 HCAPLUS
- (14) Kim, J; J Bioorg Med Chem 1998, V6, P163 HCAPLUS
- (15) Kim, J; J Med Chem 1997, V40, P2818 HCAPLUS
- (16) Koo, H; Proc Natl Acad Sci 1991, V88, P1212 HCAPLUS
- (17) Liu, L; Annu Rev Biochem 1989, V58, P351 HCAPLUS
- (18) Liu, L; J Biol Chem 1983, V258, P15365 HCAPLUS
- (19) Liu, L; Proc Natl Acad Sci 1987, V84, P7024 HCAPLUS
- (20) Marky, L; J Biopolymers 1987, V26, P1601 HCAPLUS
- (21) McGhee, J; Biopolymers 1976, V15, P1345 HCAPLUS
- (22) Mosman, T; J Immunol Meth 1983, V65, P55
- (23) Niwa, H; Tett Lett 1981, V22, P4239 HCAPLUS
- (24) Phillips, M; J Chem Soc 1928, P2393 HCAPLUS
- (25) Pilch, D; Biochemistry 1995, V34, P16107 HCAPLUS
- (26) Pilch, D; J Proc Natl Acad Sci 1997, V94, P13565 HCAPLUS
- (27) Pommier, Y; Biochim Biophys Acta 1998, V1400, P83 HCAPLUS

(28) Preston, P; Benzimidazoles Benzimidazoles and Congeneric Tricyclic Compounds Part 1 1981, P1 HCAPLUS
 (29) Redinbo, M; Science 1998, V279, P1504 HCAPLUS
 (30) Rubin, E; J Biol Chem 1994, V269, P2433 HCAPLUS
 (31) Sun, Q; J Med Chem 1995, V38, P3638 HCAPLUS
 (32) Svejstrup, J; J Mol Biol 1991, V222, P669 HCAPLUS
 (33) Tamura, H; Nuclei Acids Res 1991, V19, P69 HCAPLUS
 (34) Tsao, Y; Cell 1989, V56, P111 HCAPLUS
 (35) Wang, J; Annu Rev Biochem 1996, V65, P635 HCAPLUS
 (36) Wells, R; J Mol Biol 1970, V54, P465 HCAPLUS
 (37) Xu, Z; Biochemistry 1998, V37, P3558 HCAPLUS
 IT 237429-44-6P
 RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses)
 (preparation of phenylterbenzimidazoles as topoisomerase I poisons)
 RN 237429-44-6 HCAPLUS
 CN [2,5':2',5''-Ter-1H-benzimidazol]-2''-amine, 5-phenyl- (9CI) (CA INDEX NAME)



L25 ANSWER 5 OF 16 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 2000:269111 HCAPLUS
 DN 133:53166
 ED Entered STN: 26 Apr 2000
 TI Heterocyclic bibenzimidazole derivatives as topoisomerase I inhibitors
 AU Jin, Song; Kim, Jung Sun; Sim, Sai-Peng; Liu, Angela; Pilch, Daniel S.; Liu, Leroy F.; LaVoie, Edmond J.
 CS Department of Pharmaceutical Chemistry, Rutgers, The State University of New Jersey, Piscataway, NJ, 08854-8020, USA
 SO Bioorganic & Medicinal Chemistry Letters (2000), 10(8), 719-723
 CODEN: BMCLE8; ISSN: 0960-894X
 PB Elsevier Science Ltd.
 DT Journal
 LA English
 CC 1-3 (Pharmacology)
 Section cross-reference(s): 7, 28
 AB A series of 2'-heterocyclic derivs. of 5-phenyl-2,5'-1H-bibenzimidazoles were evaluated for topoisomerase I poisoning activity and cytotoxicity. Topo I poisoning activity was associated with 2'-derivs. that possessed a hydrogen atom capable of hydrogen bond formation, suggesting that the interat. distances between such hydrogen atoms and the heteroatoms on the adjacent benzimidazole influence activity.
 ST heterocyclic bibenzimidazole prepn topoisomerase I inhibitor
 IT Molecular modeling
 Structure-activity relationship
 (preparation of heterocyclic bibenzimidazole derivs. as topoisomerase I inhibitors)
 IT Cytotoxic agents
 (preparation of heterocyclic bibenzimidazole derivs. as topoisomerase I inhibitors and cytotoxic agents)
 IT Proliferation inhibition
 (proliferation inhibitors; preparation of heterocyclic bibenzimidazole derivs. as topoisomerase I inhibitors and cytotoxic agents)
 IT 167959-27-5P 230308-96-0P 230308-97-1P 230308-98-2P
 277754-98-0P 277754-99-1P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of heterocyclic bibenzimidazole derivs. as topoisomerase I inhibitors)
 IT 143180-75-0
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(preparation of heterocyclic bibenzimidazole derivs. as topoisomerase I inhibitors)

IT 1196-69-6, 5-Formylindole 1196-70-9, 6-Formylindole 230308-95-9
237429-60-6

RL: RCT (Reactant); RACT (Reactant or reagent)
(preparation of heterocyclic bibenzimidazole derivs. as topoisomerase I inhibitors)

RE.CNT 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE


(1) Beerman, T; Biochem Biophys Acta 1992, V1131, P53 HCAPLUS
(2) Chen, A; Cancer Res 1993, V53, P1332 HCAPLUS
(3) Chen, A; Proc Natl Acad Sci USA 1993, V90, P8131 HCAPLUS
(4) Jones, R; Org Synth 1950, V30, P86 HCAPLUS
(5) Kim, J; J Med Chem 1997, V40, P2818 HCAPLUS
(6) Moyer, M; J Org Chem 1986, V51, P5106 HCAPLUS
(7) Ohmori, J; J Med Chem 1996, V39, P1331 HCAPLUS
(8) Pilch, D; Proc Natl Acad Sci USA 1997, V94, P13565 HCAPLUS
(9) Rangarajan, M; to be published in Bioorg Med Chem
(10) Rewcastle, G; J Med Chem 1996, V39, P918
(11) Sun, Q; J Med Chem 1995, V38, P3638 HCAPLUS
(12) Xu, Z; Biochemistry 1998, V37, P3558 HCAPLUS

IT 167959-27-5P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of heterocyclic bibenzimidazole derivs. as topoisomerase I inhibitors)

RN 167959-27-5 HCAPLUS

CN 2,5':2',5''-Ter-1H-benzimidazole, 5-phenyl- (9CI) (CA INDEX NAME)



L25 ANSWER 6 OF 16 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1999:529136 HCAPLUS

DN 131:157763

ED Entered STN: 24 Aug 1999

TI Preparation of terbenzimidazoles as topoisomerase I inhibitors

IN Lavoie, Edmond J.; Kim, Jung Sun; Rangarajan, Meera;
Liu, Leroy Fong

PA Rutgers, the State University of New Jersey, USA

SO PCT Int. Appl., 60 pp.
CODEN: PIXXD2

DT Patent

LA English

IC ICM C07D235-18
ICS C07D235-26; C07D235-28; C07D235-30; A61K031-415; C07D491-04;
C07D235-02; C07D491-04; C07D317-00; C07D235-00

CC 28-9 (Heterocyclic Compounds (More Than One Hetero Atom))
Section cross-reference(s): 1

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9941241	A1	19990819	WO 1999-US2966	19990212
W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
US 6063801	A	20000516	US 1998-23147	19980212
CA 2318347	AA	19990819	CA 1999-2318347	19990212
AU 9926724	A1	19990830	AU 1999-26724	19990212
AU 748778	B2	20020613		
EP 1054870	A1	20001129	EP 1999-906928	19990212
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,			

Search done by Noble Jarrell


IE, FI					
JP 2002503653	T2	20020205	JP 2000-531436	19990212	
US 6221892	B1	20010424	US 2000-484402	20000114	
US 2001009919	A1	20010726	US 2001-796500	20010228	
US 6399642	B2	20020604			
PRAI US 1998-23147	A2	19980212			
WO 1999-US2966	W	19990212			
US 2000-484402	A1	20000114			

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
WO 9941241	ICM ICS	C07D235-18 C07D235-26; C07D235-28; C07D235-30; A61K031-415; C07D491-04; C07D235-02; C07D491-04; C07D317-00; C07D235-00
US 6063801	ECLA	C07D235/18; C07D235/26; C07D235/28; C07D235/30; C07D491/04+317A+235A
US 6221892	ECLA	C07D235/18; C07D235/26; C07D235/28; C07D235/30; C07D491/04+317A+235A
US 2001009919	ECLA	C07D235/18; C07D235/26; C07D235/28; C07D235/30; C07D491/04+317A+235A

OS MARPAT 131:157763

GI



AB Title compds. [I; R1,R2 = H, halo, alkyl, alkoxy, etc.; R3-R5 = H, alkoxycarbonyl, (hetero)aryl(alkyl), etc.; R6,R7 = H, halo, alkyl, alkoxy, etc.; R8 = halo, (halo)alkyl, CO₂H, etc.] were prepared Thus, 5-phenyl-2-(3,4-diaminophenyl)benzimidazole was cyclocondensed with 5-formyl-2-trifluoromethylbenzimidazole (preparation each given) to give I (R1,R3-R7 = H, R2 = Ph, R8 = CF₃). Data for biol. activity of I were given.

ST terbenzimidazole prepn topoisomerase I inhibitor; antitumor

IT terbenzimidazole prepn; antifungal terbenzimidazole prepn

IT Antitumor agents

Fungicides

(preparation of terbenzimidazoles as topoisomerase I inhibitors)

IT 143180-75-0

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(mediated disorders; treatment; preparation of terbenzimidazoles as topoisomerase I inhibitors)

IT 167959-27-5P 185199-36-4P 192879-73-5P

192879-74-6P 192879-75-7P 237429-42-4P

237429-43-5P 237429-44-6P 237429-45-7P

237429-46-8P 237429-47-9P 237429-48-0P

237429-49-1P 237429-50-4P 237429-51-5P

237429-52-6P 237429-53-7P 237429-54-8P

237429-55-9P 237429-56-0P 237429-57-1P

237429-58-2P 237429-59-3P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of terbenzimidazoles as topoisomerase I inhibitors)

IT 98-80-6, Phenylboronic acid 107-95-9, .beta.-Alanine 495-69-2,

Hippuric acid 591-19-5, 3-Bromoaniline 875-51-4, 4-Bromo-2-

nitroaniline 2544-06-1, 3-Methoxypropionic acid 6393-40-4,

4-Amino-3-nitrobenzonitrile 6943-69-7 31433-98-4 35998-98-2,

3,4-Dinitrobenzaldehyde 167959-20-8 192879-70-2 192879-72-4

RL: RCT (Reactant); RACT (Reactant or reagent)

(preparation of terbenzimidazoles as topoisomerase I inhibitors)

IT 615-55-4P, 3,4-Dibromoaniline 1575-37-7P, 4-Bromo-1,2-diaminobenzene

Search done by Noble Jarrell

3671-61-2P 17626-40-3P, 3,4-Diaminobenzonitrile 24108-97-2P,
 N-(3,4-Dibromophenyl)acetamide 49764-63-8P, 4,5-Dibromo-1,2-
 diaminobenzene 75293-96-8P 75293-97-9P, 3,4-Dibromo-6-nitroaniline
 106429-45-2P 106429-59-8P 108447-01-4P 117878-22-5P,
 [1,1':2',1''-Terphenyl]-4',5'-diamine 185199-45-5P 221289-88-9P
 230308-95-9P 237429-60-6P 237429-61-7P 237429-62-8P 237429-63-9P
 237429-64-0P 237429-65-1P 237429-66-2P 237429-67-3P 237429-68-4P
 237429-69-5P 237429-70-8P 237429-71-9P 237429-72-0P 237429-73-1P
 237429-74-2P 237429-75-3P 237429-76-4P 237429-77-5P 237429-78-6P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
 (Reactant or reagent)

(preparation of terbenzimidazoles as topoisomerase I inhibitors)

RE.CNT 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Chen, A; Cancer Research 1993, V53(6), P1332 HCAPLUS
- (2) Goldman, G; Biochemistry 1997, V36(21), P6488 HCAPLUS
- (3) Kim, J; Journal of Medicinal Chemistry 1997, V40(18), P2818 HCAPLUS
- (4) Pilch, D; Proceedings of the National Academy of Sciences of USA 1997,
 V94(2), P13565
- (5) Rutgers, The State University of New Jersey; WO 9636612 A 1996 HCAPLUS
- (6) Rutgers, The State University of New Jersey; WO 9831673 A 1998 HCAPLUS
- (7) Sun, Q; Journal of Medicinal Chemistry 1995, V38(18), P3638 HCAPLUS


IT 167959-27-5P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological
 study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use);
 BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of terbenzimidazoles as topoisomerase I inhibitors)

RN 167959-27-5 HCAPLUS

CN 2,5':2',5''-Ter-1H-benzimidazole, 5-phenyl- (9CI) (CA INDEX NAME)



L25 ANSWER 7 OF 16 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1999:460416 HCAPLUS

DN 131:87914

ED Entered STN: 28 Jul 1999

TI Heterocyclic topoisomerase poisons, namely 2-(benzimidazol-5-
 yl)benzimidazoles

IN Lavoie, Edmond J.; Kim, Jun Sung; Liu, Leroy Fong

PA Rutgers, the State University of New Jersey, USA

SO PCT Int. Appl., 29 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM C07D403-04

ICS A61K031-415

CC 28-9 (Heterocyclic Compounds (More Than One Hetero Atom))

Section cross-reference(s): 1, 7

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9933824	A1	19990708	WO 1998-US27822	19981230
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
CA 2316221	AA	19990708	CA 1998-2316221	19981230
AU 9920220	A1	19990719	AU 1999-20220	19981230
AU 753268	B2	20021010		
EP 1044199	A1	20001018	EP 1998-965021	19981230
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
JP 2002509858	T2	20020402	JP 2000-526506	19981230

Search done by Noble Jarrell


US 2002151575	A1	20021017	US 2001-869141	20010613
US 6667302	B2	20031223		
US 2004082637	A1	20040429	US 2003-690800	20031021
PRAI US 1997-70287P	P	19971231		
WO 1998-US27822	W	19981230		
US 2001-869141	A3	20010613		

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
WO 9933824	ICM	C07D403-04
	ICS	A61K031-415
WO 9933824	ECLA	C07D235/20; C07D403/14+239+235C+235C; C07D403/14+249+235C+235C
US 2002151575	ECLA	C07D235/20; C07D403/14+239+235C+235C; C07D403/14+249+235C+235C
US 2004082637	ECLA	C07D235/20; C07D403/14+239+235C+235C; C07D403/14+249+235C+235C

OS MARPAT 131:87914

GI



AB The invention provides title compds. I [R₁, R₂ = H, alkyl, cycloalkyl, alkoxy, (un)substituted (hetero)aryl, etc; or R₁R₂ = benzo, methylenedioxy; R₃ = H, alkyl, cycloalkyl, alkoxy, OH, CF₃O, halo, etc.; R₄R₅ = 3- to 5-atom ring-forming chain containing .gtoreq.1 NH group, and as further units O (except peroxides), S, N(X), C, or C(O), where X = null, H, O, alkyl, Ph, or PhCH₂], as well as their pharmaceutically acceptable salts, pharmaceutical compns., and use of any of these to treat cancer. For instance, 5-phenyl-2-[2-(3,4-dinitrophenyl)benzimidazol-5-yl]benzimidazole was hydrogenated over Pd/C to give the 3,4-diamino compound, which underwent diazotization with concomitant cyclization to give title compound II. Two example compds. potently inhibited topoisomerase I in vitro, and also exhibited cytotoxic activity against RPMI 8402 cancer cells and camptothecin-resistant CPT-K5 cells in vitro.

ST heterocyclic topoisomerase poison benzimidazolylbenzimidazole prepn

IT Antitumor agents

Cytotoxic agents

(preparation of (benzimidazolyl)benzimidazoles as topoisomerase poisons for use as anticancer agents)

IT Antitumor agents

(solid tumor, treatment; preparation of (benzimidazolyl)benzimidazoles as topoisomerase poisons for use as anticancer agents)

IT 167959-27-5, 5-Phenyl-2-[2-(benzimidazol-5-yl)benzimidazol-5-yl]benzimidazole

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)


(comparison compound; preparation of (benzimidazolyl)benzimidazoles as topoisomerase poisons for use as anticancer agents)

IT 230308-98-2P, 5-Phenyl-2-[2-(quinoxalin-6-yl)benzimidazol-5-yl]benzimidazole

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(comparison compound; preparation of (benzimidazolyl)benzimidazoles as

topoisomerase poisons for use as anticancer agents)
 IT 230308-95-9P, 5-Phenyl-2-[2-(3,4-diaminophenyl)benzimidazol-5-yl]benzimidazole
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (intermediate; preparation of (benzimidazolyl)benzimidazoles as topoisomerase poisons for use as anticancer agents)
 IT 143180-75-0
 RL: BPR (Biological process); BSU (Biological study, unclassified); MSC (Miscellaneous); BIOL (Biological study); PROC (Process)
 (preparation of (benzimidazolyl)benzimidazoles as topoisomerase poisons for use as anticancer agents)
 IT 144-62-7, Ethanedioic acid, reactions 517-21-5, Glyoxal disodium bisulfite 192879-72-4, 5-Phenyl-2-[2-(3,4-dinitrophenyl)benzimidazol-5-yl]benzimidazole
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (starting material; preparation of (benzimidazolyl)benzimidazoles as topoisomerase poisons for use as anticancer agents)
 IT 230308-96-0P, 5-Phenyl-2-[2-(1H-benzotriazol-5-yl)benzimidazol-5-yl]benzimidazole 230308-97-1P, 5-Phenyl-2-[2-(1,2,3,4-tetrahydro-2,3-dioxoquinoxalin-6-yl)benzimidazol-5-yl]benzimidazole
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (target compound; preparation of (benzimidazolyl)benzimidazoles as topoisomerase poisons for use as anticancer agents)
 RE.CNT 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD
 RE
 (1) Arznad; ARZNEIM-FORSCH 1974, V24(12), P1927
 (2) Goldman, G; BIOCHEMISTRY 1997, V36(21), P6488 HCAPLUS
 (3) Heinz, L; US 3538097 A 1970
 (4) Kim, J; J MED CHEM 1996, V39(4), P992 HCAPLUS
 (5) Kim, J; JOURNAL OF MEDICINAL CHEMISTRY 1997, V40(18), P2818 HCAPLUS
 (6) Lavoie, E; WO 9636612 A 1996 HCAPLUS
 (7) Lavoie, E; WO 9831673 A 1998 HCAPLUS
 (8) Loewe, H; Basic substituted 2,6-bisbenzimidazole derivatives, a novel series of substances with chemotherapeutic activity 1975, 17, HCAPLUS
 (9) Sun, Q; BIOORGANIC & MEDICINAL CHEMISTRY LETTERS 1994, V4(24), P2871 HCAPLUS
 IT 167959-27-5, 5-Phenyl-2-[2-(benzimidazol-5-yl)benzimidazol-5-yl]benzimidazole
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)
 (comparison compound; preparation of (benzimidazolyl)benzimidazoles as topoisomerase poisons for use as anticancer agents)
 RN 167959-27-5 HCAPLUS
 CN 2,5':2',5''-Ter-1H-benzimidazole, 5-phenyl- (9CI) (CA INDEX NAME)




L25 ANSWER 8 OF 16 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 1998:427779 HCAPLUS
 DN 129:62952
 ED Entered STN: 11 Jul 1998
 TI Terbenzimidazoles useful as antifungal agents
 IN Lavoie, Edmond J.; Liu, Leroy Fong; Sun, Qun
 PA Rutgers, the State University of New Jersey, USA
 SO U.S., 22 pp., Cont.-in-part of U.S. Ser. No. 618,988.
 CODEN: USXXAM
 DT Patent
 LA English
 IC ICM A61K031-415
 NCL 514394000
 CC 1-5 (Pharmacology)
 Section cross-reference(s): 28
 FAN.CNT 2

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----

PI US 5770617 A 19980623 US 1997-786629 19970121
 US 5767142 A 19980616 US 1996-618988 19960320
 CA 2278452 AA 19980723 CA 1998-2278452 19980121
 WO 9831673 A1 19980723 WO 1998-US1005 19980121
 W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE,
 DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG,
 KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX,
 NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT,
 UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
 RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI,
 FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM,
 GA, GN, ML, MR, NE, SN, TD, TG
 AU 9861327 A1 19980807 AU 1998-61327 19980121
 AU 746663 B2 20020502
 EP 960103 A1 19991201 EP 1998-905972 19980121
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
 IE, FI
 BR 9806923 A 20000418 BR 1998-6923 19980121
 NZ 336606 A 20010427 NZ 1998-336606 19980121
 JP 2002513399 T2 20020508 JP 1998-534624 19980121
 MX 9906793 A 20000228 MX 1999-6793 19990721
 AU 9952697 A1 19991202 AU 1999-52697 19991005
 AU 730456 B2 20010308
 PRAI US 1996-618988 A2 19960320
 AU 1996-57466 A3 19960514
 US 1997-786629 A 19970121
 WO 1998-US1005 W 19980121

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
US 5770617	ICM	A61K031-415
	NCL	514394000
US 5767142	ECLA	C07D235/18; C07D401/14+235C+235C+235C+213; C07D401/14R+235C+213
WO 9831673	ECLA	C07D235/18; C07D401/14R+235C+213
OS MARPAT 129:62952		
GI		



AB The present invention provides a method of treating fungal infection with an antifungal topoisomerase I inhibitor I [Ar = aryl, heteroaryl, benzo; X = H, CN, CHO, OH, acetyl, CF3, alkoxy, NO2, NH2, halogen, haloalkyl; Y = H, alkyl, aralkyl; Y1 = H, alkyl; n = 0, 1; Z = H, alkyl, halogen, haloalkyl] or their pharmaceutically acceptable salts. Thus, I [X = Y = Y1 = Z = H; Ar = Ph, n = 1, II] was obtained from 5-benzimidazolecarboxylic acid and 4-phenyl-1,2-phenylenediamine in 4 steps. II is about one half as potent as Hoechst 33342 as a topoisomerase I inhibitor.

ST terbenzimidazole prepn topoisomerase I inhibitor fungicide

IT Fungicides
 (terbenzimidazoles useful as antifungal agents)

IT 167959-21-9P, 2,5':2',5''-Ter-1H-benzimidazole
 167959-22-0P 167959-24-2P 167959-25-3P
 167959-26-4P 167959-27-5P 185199-36-4P
 185199-38-6P 185199-39-7P 209126-70-5P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (terbenzimidazoles useful as antifungal agents)

IT 143180-75-0
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
 (terbenzimidazoles useful as antifungal agents)

IT 88-74-4, 2-Nitroaniline 106-39-8, 4-Bromo-1-chlorobenzene 1635-61-6,
 5-Chloro-2-nitroaniline 6393-40-4, 4-Amino-3-nitrobenzonitrile
 15788-16-6, 5-Benzimidazolecarboxylic acid 17626-40-3,
 3,4-Diaminobenzonitrile 58442-17-4, 1H-Benzimidazole-5-carboxaldehyde

62579-61-7

RL: RCT (Reactant); RACT (Reactant or reagent)

(terbenzimidazoles useful as antifungal agents)

IT 95-83-0P, 4-Chloro-1,2-phenylenediamine 875-51-4P, 4-Bromo-2-nitroaniline 1575-37-7P, 4-Bromo-1,2-phenylenediamine 4085-18-1P, 4-Phenyl-2-nitroaniline 17151-48-3P 59656-62-1P 160522-85-0P 167959-13-9P 167959-18-4P 167959-19-5P 167959-20-8P 185199-45-5P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(terbenzimidazoles useful as antifungal agents)

RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

(1) Dykstra; US 5639755 1997 HCAPLUS

(2) Dykstra; US 5643935 1997 HCAPLUS


IT 167959-21-9P, 2,5':2',5''-Ter-1H-benzimidazole

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(terbenzimidazoles useful as antifungal agents)

RN 167959-21-9 HCAPLUS

CN 2,5':2',5''-Ter-1H-benzimidazole (9CI) (CA INDEX NAME)



L25 ANSWER 9 OF 16 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1998:168435 HCAPLUS

DN 128:278724

ED Entered STN: 21 Mar 1998

TI Quantitative structure-activity relationships on 5-substituted terbenzimidazoles as topoisomerase I poisons and antitumor agents

AU Kim, Jung Sun; Sun, Qun; Yu, Chiang; Liu, Angela; Liu, Leroy F.; Lavoie, Edmond J.

CS Department of Pharmaceutical Chemistry, Rutgers, The State University of New Jersey, Piscataway, NJ, 08855, USA

SO Bioorganic & Medicinal Chemistry (1998), 6(2), 163-172

CODEN: BMECEP; ISSN: 0968-0896

PB Elsevier Science Ltd.

DT Journal

LA English

CC 1-6 (Pharmacology)

Section cross-reference(s): 28

AB 5-Substituted terbenzimidazoles were synthesized and evaluated as mammalian topoisomerase I poisons and for cytotoxicity against a human lymphoblastoma cell line, RPMI-8402. No correlation was observed between topoisomerase I poisoning activity and the Hansch .pi. value or the .sigma.meta and .sigma.para values associated with each substituent. These data suggest that electronic effects and relative lipophilicity of substituents at the 5-position of these terbenzimidazoles do not have a significant effect upon intrinsic topoisomerase I poisoning activity. A good correlation between the relative .pi. values for the various substituents evaluated and cytotoxic activity was noted. Exptl. determined log P values did not correlate well with either cytotoxicity or .pi. values. Capacity factors (log k') as determined by high pressure liquid chromatog. did correlate well with the .pi. values of varied substituents and cytotoxicity. These data indicated that the relative lipophilic activity of substituents at the 5-position of these terbenzimidazoles can strongly influence relative cytotoxic activity.

ST terbenzimidazole topoisomerase I poisoning activity; cytotoxicity human lymphoblast cell; antitumor agent terbenzimidazole prepn; benzimidazole ter topoisomerase I poisoning activity

IT Antitumor agents

Cytotoxicity

QSAR (structure-activity relationship)

(QSAR on 5-substituted terbenzimidazoles as topoisomerase I poisons and antitumor agents)

IT 167959-21-9, 2,5':2',5''-Ter-1H-benzimidazole 167959-22-0

167959-23-1 167959-27-5 192879-67-7

Search done by Noble Jarrell

192879-68-8
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)
 (QSAR on 5-substituted terbenzimidazoles as topoisomerase I poisons and antitumor agents)

IT 185199-36-4P 185199-38-6P 205749-93-5P
 205749-94-6P 205749-95-7P 205749-96-8P
 205749-97-9P, [2,5':2',5''-Ter-1H-benzimidazol]-5-ol
 205749-98-0P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (QSAR on 5-substituted terbenzimidazoles as topoisomerase I poisons and antitumor agents)

IT 143180-75-0
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
 (QSAR on 5-substituted terbenzimidazoles as topoisomerase I poisons and antitumor agents)

IT 96-96-8, 2-Nitro-4-methoxyaniline 99-56-9, 4-Nitrophenylene-1,2-diamine
 364-78-3, 2-Nitro-4-fluoroaniline 875-51-4, 2-Nitro-4-bromoaniline
 1635-61-6, 2-Nitro-5-chloroaniline 54997-99-8 167959-20-8
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (QSAR on 5-substituted terbenzimidazoles as topoisomerase I poisons and antitumor agents)

IT 95-83-0P 102-51-2P 367-31-7P 1575-37-7P 155198-10-0P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (QSAR on 5-substituted terbenzimidazoles as topoisomerase I poisons and antitumor agents)

RE.CNT 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD


RE

- (1) Beerman, T; Biochim Biophys Acta 1992, V1131, P53 HCAPLUS
- (2) Bodley, A; Bio/technology 1988, V6, P1315 HCAPLUS
- (3) Carmichael, J; Cancer Res 1987, V47, P936 HCAPLUS
- (4) Chen, A; Annu Rev Pharmacol Toxicol 1994, V34, P191 HCAPLUS
- (5) Chen, A; Cancer Res 1993, V53, P1332 HCAPLUS
- (6) Chen, A; Proc Natl Acad Sci USA 1993, V90, P8131 HCAPLUS
- (7) Coburn, R; QSAR-PC: PAR 1987
- (8) Craig, P; J Med Chem 1971, V14, P680 HCAPLUS
- (9) Dearden, J; Quant Struct-Act Relat 1988, V7, P133 HCAPLUS
- (10) Denizot, F; J Immunol Methods 1986, V89, P271 MEDLINE
- (11) D'Arpa, P; Biochim Biophys Acta 1989, V989, P163 HCAPLUS
- (12) Gatto, B; Cancer Res 1996, V56, P2759
- (13) Hansch, C; Exploring QSAR-hydrophobic, electronic, and steric constraints 1995
- (14) Kim, J; J Med Chem 1996, V39, P992 HCAPLUS
- (15) Kim, J; J Med Chem 1997, V40, P2818 HCAPLUS
- (16) Liu, L; Annu Rev Biochem 1989, V58, P351 HCAPLUS
- (17) Liu, L; J Biol Chem 1983, V258, P15365 HCAPLUS
- (18) Loewe, H; Arzneim-Forsch 1974, V24, P1927 HCAPLUS
- (19) Maniatis, T; Molecular Cloning, a Laboratory Manual 1982, P149
- (20) Minick, D; J Liq Chromatogr 1987, V10, P2565 HCAPLUS
- (21) Mosmann, T; J Immunol Methods 1983, V65, P55 MEDLINE
- (22) Nowotnik, D; J Chromatogr 1993, V630, P105
- (23) Rekker, R; Calculations of drug lipophilicity 1992
- (24) Schneider, E; Adv in Pharmacol 1990, V21, P149 HCAPLUS
- (25) Sun, Q; Bioorg Med Chem Lett 1994, V4, P2871 HCAPLUS
- (26) Sun, Q; J Med Chem 1995, V38, P3638 HCAPLUS
- (27) Wang, J; Annu Rev Biochem 1985, V54, P665 MEDLINE
- (28) Wang, J; Annu Rev of Biochem 1996, V65, P635 HCAPLUS

IT 167959-21-9, 2,5':2',5''-Ter-1H-benzimidazole
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)
 (QSAR on 5-substituted terbenzimidazoles as topoisomerase I poisons and antitumor agents)


RN 167959-21-9 HCAPLUS

CN 2,5':2',5''-Ter-1H-benzimidazole (9CI) (CA INDEX NAME)



L25 ANSWER 10 OF 16 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 1998:113031 HCAPLUS
 DN 128:239061
 ED Entered STN: 26 Feb 1998
 TI DNA Minor Groove Binding-Directed Poisoning of Human DNA Topoisomerase I
 by Terbenzimidazoles
 AU Xu, Zhitao; Li, Tsai-Kun; Kim, Jung Sun; LaVoie, Edmond
 J.; Breslauer, Kenneth J.; Liu, Leroy F.; Pilch, Daniel S.
 CS Department of Pharmacology, University of Medicine and Dentistry of New
 Jersey Robert Wood Johnson Medical School, Piscataway, NJ, 08854, USA
 SO Biochemistry (1998), 37(10), 3558-3566
 CODEN: BICHAW; ISSN: 0006-2960
 PB American Chemical Society
 DT Journal
 LA English
 CC 1-6 (Pharmacology)
 AB We have employed a broad range of spectroscopic, calorimetric, DNA
 cleavage, and DNA winding/unwinding measurements to characterize the DNA
 binding and topoisomerase I (TOPI) poisoning properties of three
 terbenzimidazole analogs, 5-phenylterbenzimidazole (5PTB),
 terbenzimidazole (TB), and 5-(naphthyl[2,3-d]imidazo-2-yl)bibenzenzimidazole
 (5NIBB), which differ with respect to the substitutions at their C5 and/or
 C6 positions. Our results reveal the following significant features. (i)
 The overall extent to which the three terbenzimidazole analogs poison
 human TOPI follows the hierarchy 5PTB > TB > 5NIBB. (ii) The impact
 of the three terbenzimidazole analogs on the superhelical state of plasmid
 DNA depends on the [total ligand] to [base pair] ratio (rbp), having no
 effect on DNA superhelicity at rbp ratios < 0.1, while weakly
 unwinding DNA at rbp ratios > 0.1. This weak DNA unwinding activity
 exhibited by the three terbenzimidazoles does not appear to be correlated
 with the abilities of these compounds to poison TOPI. (iii) Upon
 complexation with both poly(dA).poly(dT) and salmon testes DNA, the
 three terbenzimidazole analogs exhibit flow linear dichroism properties
 characteristic of a minor groove-directed mode of binding to these host
 DNA duplexes. (iv) The apparent minor groove binding affinities of the
 three terbenzimidazole analogs for the d(GA4T4C)2 duplex follow a qual.
 similar hierarchy to that noted above for ligand-induced poisoning of
 human TOPI-namely, 5PTB > TB > 5NIBB. In the aggregate, our results
 suggest that DNA minor groove binding, but not DNA unwinding, is important
 in the poisoning of TOPI by terbenzimidazoles.
 ST DAN topoisomerase I poison terbenzimidazole
 IT DNA
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL
 (Biological study); PROC (Process)
 (DNA minor groove binding in poisoning of topoisomerase I by
 terbenzimidazoles)
 IT 167959-21-9, Terbenzimidazole 167959-27-5
 192879-63-3
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological
 study, unclassified); BIOL (Biological study)
 (DNA minor groove binding in poisoning of topoisomerase I by
 terbenzimidazoles)
 IT 143180-75-0
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL
 (Biological study); PROC (Process)
 (DNA minor groove binding in poisoning of topoisomerase I by
 terbenzimidazoles)
 RE.CNT 44 THERE ARE 44 CITED REFERENCES AVAILABLE FOR THIS RECORD
 RE
 (1) Bodley, A; Cancer Res 1989, V49, P5969 HCAPLUS
 (2) Breslauer, K; Methods Enzymol 1992, V211, P533 HCAPLUS
 (3) Caserta, M; Nucleic Acids Res 1989, V17, P8463 HCAPLUS
 (4) Chen, A; Cancer Res 1993, V53, P1332 HCAPLUS
 (5) Chen, A; Proc Natl Acad Sci U S A 1993, V90, P8131 HCAPLUS
 (6) Chou, W; J Biomol Struct Dyn 1987, V5, P345 HCAPLUS

- (7) Christianson, T; Gene 1992, V110, P119 HCAPLUS
 (8) Clark, G; Biochemistry 1996, V35, P13745 HCAPLUS
 (9) Crothers, D; Biopolymers 1971, V10, P2147 HCAPLUS
 (10) Fede, A; Biochemistry 1991, V30, P11377 HCAPLUS
 (11) Fede, A; Structure 1993, V1, P177 HCAPLUS
 (12) Gatto, B; Cancer Res 1996, V56, P2795 HCAPLUS
 (13) Geacintov, N; Biochemistry 1978, V17, P5256 HCAPLUS
 (14) Geacintov, N; Biochemistry 1987, V26, P3087 HCAPLUS
 (15) Griswold, B; Anal Chem 1951, V23, P192 HCAPLUS
 (16) Gulbransen, E; J Am Chem Soc 1934, V56, P2637 HCAPLUS
 (17) Hertzberg, R; Biochemistry 1989, V28, P4629 HCAPLUS
 (18) Hertzberg, R; J Biol Chem 1990, V265, P19287 HCAPLUS
 (19) Hsiang, Y; J Biol Chem 1985, V260, P14873 HCAPLUS
 (20) Kim, J; J Med Chem 1996, V39, P992 HCAPLUS
 (21) Krogh, S; Nucleic Acids Res 1991, V19, P1235 HCAPLUS
 (22) Marky, L; Biopolymers 1987, V26, P1601 HCAPLUS
 (23) Mudd, C; J Biochem Biophys Methods 1988, V17, P171 HCAPLUS
 (24) Norden, B; Appl Spectrosc Rev 1978, V14, P157 HCAPLUS
 (25) Norden, B; Q Rev Biophys 1992, V25, P51 HCAPLUS
 (26) Parkinson, J; Biochemistry 1990, V29, P10181 HCAPLUS
 (27) Pilch, D; Biochemistry 1995, V34, P9962 HCAPLUS
 (28) Pilch, D; Biochemistry 1997, V36, P12542 HCAPLUS
 (29) Pilch, D; Drug Des Discuss 1996, V13(3-4), P115 HCAPLUS
 (30) Pilch, D; Proc Natl Acad Sci U S A 1996, V93, P8306 HCAPLUS
 (31) Pilch, D; Proc Natl Acad Sci U S A 1997, V94, P13565 HCAPLUS
 (32) Pjura, P; J Mol Biol 1987, V197, P257 HCAPLUS
 (33) Pommier, Y; Proc Natl Acad Sci U S A 1995, V92, P8861 HCAPLUS
 (34) Porter, S; Nucleic Acids Res 1989, V17, P8521 HCAPLUS
 (35) Potmesil, M; Camptothecins: New Anticancer Agents 1995
 (36) Remeta, D; Biochemistry 1991, V30, P9799 HCAPLUS
 (37) Robinson, A; J Am Chem Soc 1932, V54, P1311 HCAPLUS
 (38) Snyder, J; Proc Natl Acad Sci U S A 1989, V86, P3968 HCAPLUS
 (39) Spink, N; Nucleic Acids Res 1994, V22, P1607 HCAPLUS
 (40) Sun, Q; Bioorg Med Chem Lett 1994, V4, P2871 HCAPLUS
 (41) Sun, Q; J Med Chem 1995, V38, P3638 HCAPLUS
 (42) Teng, M; Nucleic Acids Res 1988, V16, P2671 HCAPLUS
 (43) Vega, M; Eur J Biochem 1994, V222, P721 HCAPLUS
 (44) Wilson, R; Biopolymers 1978, V17, P1235 HCAPLUS
 IT 167959-21-9, Terbenzimidazole
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)
 (DNA minor groove binding in poisoning of topoisomerase I by terbenzimidazoles)
 RN 167959-21-9 HCAPLUS
 CN 2,5':2',5''-Ter-1H-benzimidazole (9CI) (CA INDEX NAME)



- L25 ANSWER 11 OF 16 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 1997:802609 HCAPLUS
 DN 128:123523
 ED Entered STN: 25 Dec 1997
 TI A terbenzimidazole that preferentially binds and conformationally alters structurally distinct DNA duplex domains: a potential mechanism for topoisomerase I poisoning
 AU Pilch, Daniel S.; Xu, Zhitao; Sun, Qun; LaVoie, Edmond J.; Liu, Leroy F.; Breslauer, Kenneth J.
 CS Dep. Pharmacol., Univ. Med. Dentistry New Jersey, Piscataway, NJ, 08854, USA
 SO Proceedings of the National Academy of Sciences of the United States of America (1997), 94(25), 13565-13570
 CODEN: PNASA6; ISSN: 0027-8424
 PB National Academy of Sciences
 DT Journal
 LA English
 CC 1-6 (Pharmacology)
 AB The terbenzimidazoles are a class of synthetic ligands that poison the

human topoisomerase I (TOP1) enzyme and promote cancer cell death. It has been proposed that drugs of this class act as TOP1 poisons by binding to the minor groove of the DNA substrate of TOP1 and altering its structure in a manner that results in enzyme-mediated DNA cleavage. To test this hypothesis, we characterize and compare the binding properties of a 5-phenylterbenzimidazole derivative (5PTB) to the d(GA4T4C)2 duplexes. The d(GA4T4C)2 duplex contains an uninterrupted 8-bp A.cntdot.T domain, which, on the basis of x-ray crystallog. data, should induce a highly hydrated "A-tract" conformation. This duplex also exhibits anomalously slow migration in a polyacrylamide gel, a feature characteristic of a noncanonical global conformational state frequently described as "bent." By contrast, the d(GT4A4C)2 duplex contains two 4-bp A.cntdot.T tracts separated by a TpA dinucleotide step, which should induce a less hydrated "B-like" conformation. This duplex also migrates normally in a polyacrylamide gel, a feature further characteristic of a global, canonical B-form duplex. Our data reveal that, at 20.degree., 5PTB exhibits an .apprx.2.3 kcal/mol greater affinity for the d(GA4T4C)2 duplex than for the d(GT4A4C)2 duplex. Significantly, we find this sequence/conformational binding specificity of 5PTB to be entropic in origin, an observation consistent with a greater degree of drug binding-induced dehydration of the more solvated d(GA4T4C)2 duplex. By contrast with the differential duplex affinity exhibited by 5PTB, netropsin and 4',6-diamidino-2-phenylindole (DAPI), two AT-specific minor groove binding ligands that are inactive as human TOP1 poisons, bind to both duplexes with similar affinities. The electrophoretic behaviors of the ligand-free and ligand-bound duplexes are consistent with 5PTB-induced bending and/or unwinding of both duplexes, which, for the d(GA4T4C)2 duplex, is synergistic with the endogenous sequence-directed electrophoretic properties of the ligand-free duplex state. By contrast, the binding to either duplex of netropsin or DAPI induces little or no change in the electrophoretic mobilities of the duplexes. Our results demonstrate that the TOP1 poison 5PTB binds differentially to and alters the structures of the two duplexes, in contrast to netropsin and DAPI, which bind with similar affinities to the two duplexes and do not significantly alter their structures. These results are consistent with a mechanism for TOP1 poisoning in which drugs such as 5PTB differentially target conformationally distinct DNA sites and induce structural changes that promote enzyme-mediated DNA cleavage.

ST topoisomerase I poison terbenzimidazole DNA conformation

IT Conformation

(B form; terbenzimidazole that preferentially binds and conformationally alters structurally distinct DNA duplex domains and potential mechanism for topoisomerase I poisoning)

IT DNA

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(cleavage; terbenzimidazole that preferentially binds and conformationally alters structurally distinct DNA duplex domains and potential mechanism for topoisomerase I poisoning)

IT DNA

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(double-stranded; terbenzimidazole that preferentially binds and conformationally alters structurally distinct DNA duplex domains and potential mechanism for topoisomerase I poisoning)

IT Antitumor agents

Apoptosis

(terbenzimidazole that preferentially binds and conformationally alters structurally distinct DNA duplex domains and potential mechanism for topoisomerase I poisoning)

IT 122799-65-9

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(duplex; terbenzimidazole that preferentially binds and conformationally alters structurally distinct DNA duplex domains and potential mechanism for topoisomerase I poisoning)

IT 1438-30-8, Netropsin 47165-04-8, 4',6-Diamidino-2-phenylindole

167959-27-5

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)

(terbenzimidazole that preferentially binds and conformationally alters structurally distinct DNA duplex domains and potential mechanism for topoisomerase I poisoning)

IT 143180-75-0

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(terbenzimidazole that preferentially binds and conformationally alters structurally distinct DNA duplex domains and potential mechanism for topoisomerase I poisoning)

RE.CNT 45 THERE ARE 45 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Berman, H; Biophys J 1992, V63, P751 HCAPLUS
- (2) Chalikian, T; Biochemistry 1994, V33, P8629 HCAPLUS
- (3) Chen, A; Annu Rev Pharmacol Toxicol 1994, V34, P191 HCAPLUS
- (4) Chen, A; Cancer Res 1993, V53, P1332 HCAPLUS
- (5) Chen, A; Proc Natl Acad Sci USA 1993, V90, P8131 HCAPLUS
- (6) Chen, J; Nucleic Acids Res 1988, V16, P6803 HCAPLUS
- (7) Clark, G; Biochemistry 1996, V35, P13745 HCAPLUS
- (8) Crothers, D; Biopolymers 1971, V10, P2147 HCAPLUS
- (9) Dickerson, R; J Mol Biol 1996, V256, P108 HCAPLUS
- (10) Drew, H; J Mol Biol 1981, V151, P535 HCAPLUS
- (11) D'Arpa, P; Biochim Biophys Acta 1989, V989, P163 HCAPLUS
- (12) Fede, A; Biochemistry 1991, V30, P11377 HCAPLUS
- (13) Fede, A; Structure 1993, V1, P177 HCAPLUS
- (14) Goodsell, D; J Mol Biol 1994, V239, P79 HCAPLUS
- (15) Griswold, B; Anal Chem 1951, V23, P192 HCAPLUS
- (16) Gulbransen, E; J Am Chem Soc 1934, V56, P2637 HCAPLUS
- (17) Hagerman, P; Biochemistry 1985, V24, P7033 HCAPLUS
- (18) Hagerman, P; Nature (London) 1986, V321, P449 HCAPLUS
- (19) Kim, J; Bioorg Med Chem 1996, V4, P621 HCAPLUS
- (20) Kim, J; J Med Chem 1996, V39, P992 HCAPLUS
- (21) Koo, H; Biochemistry 1990, V29, P4227 HCAPLUS
- (22) Koo, H; Nature (London) 1986, V320, P501 HCAPLUS
- (23) Kopka, M; J Mol Biol 1985, V183, P553 HCAPLUS
- (24) Kopka, M; Proc Natl Acad Sci USA 1985, V82, P1376 HCAPLUS
- (25) Larsen, T; J Biomol Struct Dyn 1989, V7, P477 HCAPLUS
- (26) Lerov, J; Biochemistry 1988, V27, P8894
- (27) Liu, L; Annu Rev Biochem 1989, V58, P351 HCAPLUS
- (28) Liu, L; DNA Topoisomerases: Topoisomerase-Targeting Drugs Advances in Pharmacology 1994, V29B
- (29) Marini, J; Proc Natl Acad Sci USA 1982, V79, P7664 HCAPLUS
- (30) Marky, L; Proc Natl Acad Sci USA 1987, V84, P4359 HCAPLUS
- (31) Mudd, C; J Biochem Biophys Methods 1988, V17, P171 HCAPLUS
- (32) Nadeau, J; Proc Natl Acad Sci USA 1989, V86, P2622 HCAPLUS
- (33) Park, Y; Proc Natl Acad Sci USA 1991, V88, P1551 HCAPLUS
- (34) Parkinson, J; Biochemistry 1990, V29, P10181 HCAPLUS
- (35) Pilch, D; Drug Des Discovery 1996, V13(3-4), P115 HCAPLUS
- (36) Pjura, P; J Mol Biol 1987, V197, P257 HCAPLUS
- (37) Remeta, D; Biochemistry 1991, V30, P9799 HCAPLUS
- (38) Robinson, A; J Am Chem Soc 1932, V54, P1311 HCAPLUS
- (39) Schneider, E; Adv Pharmacol 1990, V21, P149 HCAPLUS
- (40) Spink, N; Nucleic Acids Res 1994, V22, P1607 HCAPLUS
- (41) Sun, Q; Bioorg Med Chem Lett 1994, V4, P2871 HCAPLUS
- (42) Sun, Q; J Med Chem 1995, V38, P3638 HCAPLUS
- (43) Teng, M; Nucleic Acids Res 1988, V16, P2671 HCAPLUS
- (44) Wang, J; Annu Rev Biochem 1985, V54, P665 MEDLINE
- (45) Wu, H; Nature (London) 1984, V308, P509 HCAPLUS


IT 167959-27-5

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)

(terbenzimidazole that preferentially binds and conformationally alters structurally distinct DNA duplex domains and potential mechanism for topoisomerase I poisoning)

RN 167959-27-5 HCAPLUS

CN 2,5':2',5''-Ter-1H-benzimidazole, 5-phenyl- (9CI) (CA INDEX NAME)



L25 ANSWER 12 OF 16 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1997:526711 HCAPLUS

DN 127:117067

ED Entered STN: 18 Aug 1997

TI Terbenzimidazoles: Influence of 2''-, 4-, and 5-Substituents on Cytotoxicity and Relative Potency as Topoisomerase I Poisons

Search done by Noble Jarrell

AU Kim, Jung Sun; Yu, Chiang; Liu, Angela; Liu, Leroy F.;
LaVoie, Edmond J.

CS Department of Pharmaceutical Chemistry, Rutgers The State
University of New Jersey, Piscataway, NJ, 08855, USA

SO Journal of Medicinal Chemistry (1997), 40(18), 2818-2824
CODEN: JMCMAR; ISSN: 0022-2623

PB American Chemical Society

DT Journal

LA English

CC 1-3 (Pharmacology)

Section cross-reference(s): 28

AB Terbenzimidazoles poison the nuclear enzyme topoisomerase I and possess
significant cytotoxic activity against several human tumor cell lines.
The relative pharmacol. activity of 4,5- and 5,6-benzoterbenzimidazoles
was compared to that of 5-phenylterbenzimidazole (3).
5,6-Benzoterbenzimidazole is inactive as a topoisomerase I poison and did
not exhibit significant cytotoxic activity. In contrast,
4,5-benzoterbenzimidazole retained activity as a topoisomerase I poison
but exhibited weak cytotoxic activity relative to 3. While
5-(1-naphthyl)terbenzimidazole is less potent than 3 as a topoisomerase I
poison and cytotoxic agent, 5-(2-naphthyl)terbenzimidazole has comparable
activity to 3. The presence of a p-methoxy or p-chloro substituent on the
Ph moiety did not dramatically alter the pharmacol. activity of 3.
Several analogs of 3 were synthesized wherein the 2''-substituent varied
from Me, Et, Pr, iso-Pr, Ph to p-methoxyphenyl. Evaluation of the
intrinsic activity of these analogs as topoisomerase I poisons indicates
that topoisomerase I poisoning was not diminished by the presence of a Me,
Et, Pr, and iso-Pr substituent at the 2''-position. Among the various
2''-substituted analogs evaluated, only in the case of
2''-(p-methoxyphenyl)-5-phenylterbenzimidazole was a significant decrease
in cytotoxicity observed

ST terbenzimidazole prepn cytotoxicity topoisomerase DNA damage

IT DNA
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL
(Biological study); PROC (Process)
(damage; preparation, cytotoxicity and relative potency as topoisomerase I
poisons of terbenzimidazoles)

IT Antitumor agents
Structure-activity relationship
(preparation, cytotoxicity and relative potency as topoisomerase I poisons
of terbenzimidazoles)

IT 185199-39-7P 192879-62-2P 192879-63-3P
192879-64-4P 192879-67-7P 192879-68-8P
192879-69-9P 192879-73-5P 192879-74-6P
192879-75-7P 192879-76-8P 192879-77-9P
192879-78-0P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological
study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use);
BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation, cytotoxicity and relative potency as topoisomerase I poisons
of terbenzimidazoles)

IT 100-52-7, Benzaldehyde, biological studies 167959-27-5
RL: BAC (Biological activity or effector, except adverse); BSU (Biological
study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES
(Uses)
(preparation, cytotoxicity and relative potency as topoisomerase I poisons
of terbenzimidazoles)

IT 143180-75-0
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL
(Biological study); PROC (Process)
(preparation, cytotoxicity and relative potency as topoisomerase I poisons
of terbenzimidazoles)

IT 75-07-0, Acetaldehyde, reactions 78-84-2, Isobutyraldehyde 90-11-9,
1-Bromonaphthalene 104-92-7, p-Bromoanisole 106-39-8,
p-Bromochlorobenzene 123-11-5, p-Methoxybenzaldehyde, reactions
123-38-6, Propionaldehyde, reactions 123-72-8, Butyraldehyde 580-13-2,
2-Bromonaphthalene 771-97-1, 2,3-Naphthalenediamine 875-51-4
938-25-0, 1,2-Naphthalenediamine 972-09-8 972-11-2 4458-39-3,
[1,1'-Biphenyl]-3,4-diamine 17151-48-3 35998-98-2 70744-47-7
102877-92-9, [1,1'-Biphenyl]-2,3-diamine 167959-20-8
RL: RCT (Reactant); RACT (Reactant or reagent)
(preparation, cytotoxicity and relative potency as topoisomerase I poisons
of terbenzimidazoles)

IT 2221-02-5P 31433-98-4P 192879-65-5P 192879-66-6P 192879-70-2P
192879-72-4P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT

(Reactant or reagent)

(preparation, cytotoxicity and relative potency as topoisomerase I poisons of terbenzimidazoles)

RE.CNT 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE


- (1) Anon; Organic Syntheses 1943, VII, P438
- (2) Beerman, T; Biochim Biophys Acta 1992, V1131, P53 HCAPLUS
- (3) Bodley, A; Bio/technology 1988, V6, P1315 HCAPLUS
- (4) Carmichael, J; Cancer Res 1987, V47, P936 HCAPLUS
- (5) Chen, A; Annu Rev Pharmacol Toxicol 1994, V34, P191 HCAPLUS
- (6) Chen, A; Cancer Res 1993, V53, P1332 HCAPLUS
- (7) Chen, A; Proc Natl Acad Sci U S A 1993, V90, P8131 HCAPLUS
- (8) Clark, G; Biochemistry 1996, V35, P13745 HCAPLUS
- (9) Denizot, F; J Immunol Methods 1986, V89, P271 MEDLINE
- (10) D'Arpa, P; Biochim Biophys Acta 1989, V989, P163 HCAPLUS
- (11) Gatto, B; Cancer Res 1996, V56, P2795 HCAPLUS
- (12) Goldstein, H; Helv Chim Acta 1941, V26, P1125
- (13) Kim, J; J Med Chem 1996, V39, P992 HCAPLUS
- (14) Liu, L; Annu Rev Biochem 1989, V58, P351 HCAPLUS
- (15) Liu, L; J Biol Chem 1983, V258, P15365 HCAPLUS
- (16) Maniatis, T; Molecular Cloning, a Laboratory Manual 1982, P149
- (17) Mosmann, T; Immunol Methods 1983, V65, P55 MEDLINE
- (18) Pilch, D; Drug Des Discovery 1996, V13, P115 HCAPLUS
- (19) Sako, S; Bull Chem Soc Jpn 1934, V9, P55 HCAPLUS
- (20) Schneider, E; Adv Pharmacol 1990, V21, P149 HCAPLUS
- (21) Sun, Q; Bioorg Med Chem Lett 1994, V4, P2871 HCAPLUS
- (22) Sun, Q; J Med Chem 1995, V38, P3638 HCAPLUS
- (23) Wang, J; Annu Rev Biochem 1985, V54, P665 MEDLINE
- (24) Wang, J; Annu Rev Biochem 1996, V65, P635 HCAPLUS

IT 185199-39-7P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation, cytotoxicity and relative potency as topoisomerase I poisons of terbenzimidazoles)

RN 185199-39-7 HCAPLUS

CN 2,5'-2',5''-Ter-1H-benzimidazole, 5-(4-chlorophenyl)- (9CI) (CA INDEX NAME)



L25 ANSWER 13 OF 16 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1997:299447 HCAPLUS

DN 126:340359

ED Entered STN: 10 May 1997

TI Differential Poisoning of Human and Aspergillus nidulans DNA Topoisomerase I by Bi- and Terbenzimidazoles

AU Goldman, Gustavo H.; Yu, Chiang; Wu, Hong-Yan; Sanders, Marilyn M.; La Voie, Edmond J.; Liu, Leroy F.

CS Department of Pharmacology Robert Wood Johnson Medical School, University of Medicine and Dentistry of New Jersey, Piscataway, NJ, USA

SO Biochemistry (1997), 36(21), 6488-6494

CODEN: BICHAW; ISSN: 0006-2960

PB American Chemical Society

DT Journal

LA English

CC 7-3 (Enzymes)

Section cross-reference(s): 10

AB DNA topoisomerase I has been partially purified from *Aspergillus nidulans*. The purified enzyme is most likely the major nuclear DNA topoisomerase I on the basis of the following findings. (1) Purified DNA topoisomerase I can relax both pos. and neg. supercoiled DNA. (2) Neither an energy cofactor nor Mg(II) is required for the relaxation or the cleavage reaction of the enzyme. On the basis of a phosphate-transfer experiment, the *Aspergillus* topoisomerase I was shown to have a mol. mass (Mr) of 105 kDa. The differential sensitivity of the human and *Aspergillus* topoisomerase I was compared using a number of known human DNA topoisomerase I poisons. Like human DNA topoisomerase I, *Aspergillus* topoisomerase I is highly sensitive to the poisoning activity of camptothecin and a number of bi- and


terbenzimidazoles. However, unlike human topoisomerase I, *Aspergillus* topoisomerase I is completely resistant to monobenzimidazoles, protoberberines (e.g. coralyne), and nitidine. Cytotoxicity studies using yeast expressing human and yeast topoisomerase I cDNAs have also demonstrated a similar differential sensitivity of yeast topoisomerase I to these human topoisomerase I poisons. These results together suggest that the nuclear fungal topoisomerase I may be sufficiently different from its human counterpart to serve as a mol. target for the development of antifungal drugs.

ST *Aspergillus* human DNA topoisomerase inhibitor terbenzimidazole
 IT *Aspergillus nidulans*
 (differential poisoning of human and *Aspergillus nidulans* DNA topoisomerase I by bi- and terbenzimidazoles)
 IT Structure-activity relationship
 (enzyme-inhibiting, DNA topoisomerase I; differential poisoning of human and *Aspergillus nidulans* DNA topoisomerase I by bi- and terbenzimidazoles)
 IT 143180-75-0P, DNA topoisomerase I
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PUR (Purification or recovery); BIOL (Biological study); PREP (Preparation)
 (differential poisoning of human and *Aspergillus nidulans* DNA topoisomerase I by bi- and terbenzimidazoles)
 IT 908-54-3, Berenil 6872-57-7, Nitidine 6872-73-7, Coralyne 7689-03-4, Camptothecin 23491-52-3, Hoechst 33342 91437-87-5 96954-35-7 113551-23-8 167959-22-0 167959-27-5 180077-27-4 189953-66-0
 RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)
 (differential poisoning of human and *Aspergillus nidulans* DNA topoisomerase I by bi- and terbenzimidazoles)
 RE.CNT 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD
 RE
 (1) Andoh, T; Proc Natl Acad Sci U S A 1987, V84, P5565 HCAPLUS
 (2) Bjornsti, M; Cancer Res 1989, V49, P6318 HCAPLUS
 (3) Chen, A; Proc Natl Acad Sci U S A 1993, V90, P8131 HCAPLUS
 (4) Postel, J; Antimicrob Agents Chemother 1992, V36, P2131 HCAPLUS
 (5) Postel, J; Antimicrob Agents Chemother 1995, V39, P586 HCAPLUS
 (6) Gatto, B; Cancer Res 1996, V56, P2795 HCAPLUS
 (7) Hsiang, Y; J Biol Chem 1985, V260, P14873 HCAPLUS
 (8) Janin, Y; J Med Chem 1993, V36, P3686 HCAPLUS
 (9) Jaxel, C; J Biol Chem 1991, V266, P20418 HCAPLUS
 (10) Kim, J; Bioorg Med Chem 1996, V4, P621 HCAPLUS
 (11) Kim, J; J Med Chem 1996, V39, P992 HCAPLUS
 (12) Knab, A; J Biol Chem 1993, V268, P22322 HCAPLUS
 (13) Leteurtre, F; J Biol Chem 1994, V269, P28702 HCAPLUS
 (14) Liu, L; Annu Rev Biochem 1989, V58, P351 HCAPLUS
 (15) Liu, L; DNA Topoisomerases:topoisomerase targeting drugs, Advances in Pharmacology 1994, V29B
 (16) Liu, L; J Biol Chem 1983, V258, P15365 HCAPLUS
 (17) Liu, L; Proc Natl Acad Sci U S A 1981, V78, P3487 HCAPLUS
 (18) Makhey, D; Bioorg Med Chem Lett 1996, V4, P781 HCAPLUS
 (19) Makhey, D; Med Chem Res 1995, V5, P1 HCAPLUS
 (20) Neu, H; DNA Topoisomerases:topoisomerase targeting drugs, Advances in Pharmacology 1994, V29B, P227
 (21) Nitiss, J; DNA Topoisomerases:topoisomerase targeting drugs, Advances in Pharmacology 1994, V29B, P201 HCAPLUS
 (22) Nitiss, J; Proc Natl Acad Sci U S A 1988, V85, P7501 HCAPLUS
 (23) Poddevin, B; Mol Pharmacol 1993, V44, P767 HCAPLUS
 (24) Pommier, Y; Semin Oncol 1996, V23, P3 HCAPLUS
 (25) Porter, S; Nucleic Acids Res 1989, V17, P8521 HCAPLUS
 (26) Rowe, T; J Biol Chem 1984, V259, P9177 HCAPLUS
 (27) Shen, L; Antimicrob Agents Chemother 1992, V36, P2778 HCAPLUS
 (28) Shen, L; DNA Topoisomerases:topoisomerase targeting drugs 1994, P227 HCAPLUS
 (29) Sun, Q; Bioorg Med Chem Lett 1994, V4, P2871 HCAPLUS
 (30) Sun, Q; J Med Chem 1995, V38, P3638 HCAPLUS
 (31) Tewey, K; Science 1984, V226, P466 HCAPLUS
 (32) Wang, J; Annu Rev Biochem 1985, V54, P655
 (33) Wang, J; Annu Rev Biochem 1996, V65, P635 HCAPLUS
 (34) Wang, L; Chem Res Toxicol 1993, V6, P813 HCAPLUS
 (35) Wang, L; Chem Res Toxicol 1996, V9, P75 HCAPLUS
 (36) Yamashita, Y; Biochemistry 1991, V30, P5838 HCAPLUS
 IT 167959-22-0
 RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)

(differential poisoning of human and *Aspergillus nidulans* DNA
topoisomerase I by bi- and terbenzimidazoles)

RN 167959-22-0 HCAPLUS

CN [2,5':2',5''-Ter-1H-benzimidazole]-5-carbonitrile (9CI) (CA INDEX NAME)



L25 ANSWER 14 OF 16 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1997:41984 HCAPLUS

DN 126:59953

ED Entered STN: 20 Jan 1997

TI Preparation of tribenzimidazoles useful as topoisomerase I inhibitors.

IN Lavoie, Edmond J.; Liu, Leroy Fong; Sun, Qun

PA Rutgers, the State University of New Jersey, USA; Lavoie, Edmond

J.; Liu, Leroy Fong; Sun, Qun

SO PCT Int. Appl., 36 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM C07D235-18

ICS A61K031-415; C07D401-14

CC 28-9 (Heterocyclic Compounds (More Than One Hetero Atom))


Section cross-reference(s): 1

FAN.CNT 2

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9636612	A1	19961121	WO 1996-US6853	19960514
W: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI				
RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN				
US 5807874	A	19980915	US 1995-442657	19950517
US 5767142	A	19980616	US 1996-618988	19960320
CA 2221248	AA	19961121	CA 1996-2221248	19960514
AU 9657466	A1	19961129	AU 1996-57466	19960514
AU 713317	B2	19991125		
EP 839140	A1	19980506	EP 1996-915784	19960514
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI				
JP 11508229	T2	19990721	JP 1996-534375	19960514
BR 9608476	A	19990817	BR 1996-8476	19960514
US 5948797	A	19990907	US 1997-782064	19970113
AU 9952697	A1	19991202	AU 1999-52697	19991005
AU 730456	B2	20010308		
PRAI US 1995-442657	A	19950517		
US 1996-618988	A	19960320		
AU 1996-57466	A3	19960514		
WO 1996-US6853	W	19960514		

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
WO 9636612	ICM	C07D235-18
	ICS	A61K031-415; C07D401-14
US 5807874	ECLA	C07D235/18; C07D401/14+235C+235C+235C+213
US 5767142	ECLA	C07D235/18; C07D401/14+235C+235C+235C+213; C07D401/14R+235C+213
US 5948797	ECLA	C07D235/18; C07D401/14+235C+235C+235C+213
OS	MARPAT	126:59953
GI		



AB Title compds. [I; Ar = 0-1 (substituted) aryl, heteroaryl which may be fused to the benzo moiety; X = H, CN, CHO, OH, Ac, CF₃, alkoxy, NO₂, NH₂, halo, haloalkyl; Y = H, alkyl, aralkyl; Y₁ = H, alkyl; ; Z = H, alkyl, halo, haloalkyl], were prepared Thus, 4-(4-pyridyl)-2-nitroaniline reacted with 5-formyl-2-(benzimidazol-5-yl)benzimidazole to give 43% 5-(4-pyridyl)-2-[2-(benzimidazol-5-yl)benzimidazol-5-yl]benzimidazole. This showed IC₅₀ = 0.01 .mu.M against KBV-1 cells.

ST tribenzimidazole prepn topoisomerase inhibitor; anticancer tribenzimidazole prepn

IT Antitumor agents
(preparation of tribenzimidazoles useful as topoisomerase I inhibitors)

IT 167959-21-9P, 2,5':2',5''-Ter-1H-benzimidazole
167959-22-0P 167959-23-1P 167959-24-2P
167959-25-3P 167959-26-4P 167959-27-5P
185199-36-4P 185199-38-6P 185199-39-7P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of tribenzimidazoles useful as topoisomerase I inhibitors)

IT 143180-75-0
RL: BPR (Biological process); BSU (Biological study, unclassified); MSC (Miscellaneous); BIOL (Biological study); PROC (Process)
(preparation of tribenzimidazoles useful as topoisomerase I inhibitors)


IT 88-74-4, 2-Nitroaniline 95-54-5, 1,2-Phenylenediamine, reactions
106-39-8, 4-Bromochlorobenzene 960-16-7, Phenyltributyltin 1635-61-6
15788-16-6, 5-Benzimidazolecarboxylic acid 17626-40-3 17997-47-6,
2-Tributylstannylpyridine 24850-33-7, Allyltributyltin 59020-10-9,
3-Tributylstannylpyridine 124252-41-1, 4-Tributylstannylpyridine
RL: RCT (Reactant); RACT (Reactant or reagent)
(preparation of tribenzimidazoles useful as topoisomerase I inhibitors)

IT 95-83-0P 875-51-4P 1575-37-7P 4085-18-1P 17151-48-3P 31433-98-4P
58442-17-4P, 1H-Benzimidazole-5-carboxaldehyde 59656-62-1P
160522-85-0P 167959-13-9P 167959-18-4P 167959-19-5P 167959-20-8P
185199-45-5P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation of tribenzimidazoles useful as topoisomerase I inhibitors)

IT 167959-21-9P, 2,5':2',5''-Ter-1H-benzimidazole
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of tribenzimidazoles useful as topoisomerase I inhibitors)


RN 167959-21-9 HCAPLUS

CN 2,5':2',5''-Ter-1H-benzimidazole (9CI) (CA INDEX NAME)



L25 ANSWER 15 OF 16 HCAPLUS COPYRIGHT 2005 ACS on STN
AN 1996:405834 HCAPLUS
DN 125:157872
ED Entered STN: 13 Jul 1996
TI Synthesis and structure-activity relationships of novel mammalian DNA topoisomerase I inhibitors (Hoechst dyes, terbenzimidazoles, antitumor

drugs)
 AU Sun, Qun
 CS Rutgers, State Univ., New Brunswick, NJ, USA
 SO (1996) 173 pp. Avail.: Univ. Microfilms Int., Order No. DA9618556
 From: Diss. Abstr. Int., B 1996, 57(2), 1093
 DT Dissertation
 LA English
 CC 1-6 (Pharmacology)
 AB Unavailable
 ST DNA topoisomerase I inhibitor structure activity
 IT Dyes
 (Hoechst; synthesis and structure-activity relationships of novel
 mammalian DNA topoisomerase I inhibitors (Hoechst dyes,
 terbenzimidazoles, antitumor drugs))
 IT Molecular structure-biological activity relationship
 Neoplasm inhibitors
 (synthesis and structure-activity relationships of novel mammalian DNA
 topoisomerase I inhibitors (Hoechst dyes, terbenzimidazoles, antitumor
 drugs))
 IT 167959-21-9D, 2,5':2',5''-Ter-1H-benzimidazole, derivs.
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological
 study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES
 (Uses)
 (synthesis and structure-activity relationships of novel mammalian DNA
 topoisomerase I inhibitors (Hoechst dyes, terbenzimidazoles, antitumor
 drugs))
 IT 143180-75-0, DNA topoisomerase I
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL
 (Biological study); PROC (Process)
 (synthesis and structure-activity relationships of novel mammalian DNA
 topoisomerase I inhibitors (Hoechst dyes, terbenzimidazoles, antitumor
 drugs))
 IT 167959-21-9D, 2,5':2',5''-Ter-1H-benzimidazole, derivs.
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological
 study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES
 (Uses)
 (synthesis and structure-activity relationships of novel mammalian DNA
 topoisomerase I inhibitors (Hoechst dyes, terbenzimidazoles, antitumor
 drugs))
 RN 167959-21-9 HCAPLUS
 CN 2,5':2',5''-Ter-1H-benzimidazole (9CI) (CA INDEX NAME)



L25 ANSWER 16 OF 16 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 1995:746950 HCAPLUS
 DN 123:198740
 ED Entered STN: 19 Aug 1995
 TI Synthesis and Evaluation of Terbenzimidazoles as Topoisomerase I
 Inhibitors
 AU Sun, Qun; Gatto, Barbara; Yu, Chiang; Liu, Angela; Liu, Leroy F.
 ; LaVoie, Edmond J.
 CS Department of Pharmaceutical Chemistry, Rutgers, State
 University of New Jersey, Piscataway, NJ, 08855, USA
 SO Journal of Medicinal Chemistry (1995), 38(18), 3638-44
 CODEN: JMCMAR; ISSN: 0022-2623
 PB American Chemical Society
 DT Journal
 LA English
 CC 28-18 (Heterocyclic Compounds (More Than One Hetero Atom))
 Section cross-reference(s): 1
 OS CASREACT 123:198740
 AB The synthesis and pharmacol. activity of a series of terbenzimidazoles are
 described. The ability of these derivs. to induce DNA cleavage in the
 presence of topoisomerase I was evaluated in vitro. These analogs were
 also assayed for their cytotoxicity in RPMI cells and the
 camptothecin-resistant CPT-K5 cells. In addition the potential for these

compds. to serve as substrates for MDR1 was also determined. Several terbenzimidazoles exhibited similar cytotoxicity against variants of human tumor cells that either overexpress MDR1 or are camptothecin-resistant. Cyclocondensation of 1,2-benzenediamine with 2,5'-bi-1H-benzimidazole-5-carboxaldehyde gave 2,5':2',5''-Ter-1H-benzimidazole.

ST terbenzimidazole prepn topoisomerase inhibitor

IT 167959-13-9P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); RCT (Reactant); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent)
(2,5'-bi-1H-benzimidazole-5-carbonitrile; preparation of terbenzimidazoles as topoisomerase I inhibitors)

IT 167959-20-8P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(2,5'-bi-1H-benzimidazole-5-carboxaldehyde; preparation of terbenzimidazoles as topoisomerase I inhibitors)

IT 167959-21-9P, 2,5':2',5''-Ter-1H-benzimidazole

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
(2,5':2',5''-Ter-1H-benzimidazole; preparation of terbenzimidazoles as topoisomerase I inhibitors)

IT 167959-14-0P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(2-(4-methoxyphenyl)-1H-Benzimidazole-5-carbonitrile; preparation of terbenzimidazoles as topoisomerase I inhibitors)

IT 167959-18-4P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(2-nitro-4-(2-pyridinyl)benzenamine; preparation of terbenzimidazoles as topoisomerase I inhibitors)

IT 167959-19-5P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(2-nitro-4-(3-pyridinyl)benzenamine; preparation of terbenzimidazoles as topoisomerase I inhibitors)

IT 143180-75-0

RL: ADV (Adverse effect, including toxicity); BIOL (Biological study)
(preparation of terbenzimidazoles as topoisomerase I inhibitors)

IT 167959-15-1P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); RCT (Reactant); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent)
(preparation of terbenzimidazoles as topoisomerase I inhibitors)

IT 23491-52-3DP, Hoechst 33342, analogs and derivs 167959-17-3P

167959-22-0P 167959-23-1P 167959-24-2P

167959-25-3P 167959-26-4P 167959-27-5P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
(preparation of terbenzimidazoles as topoisomerase I inhibitors)

IT 88-74-4, Benzenamine, 2-nitro 95-54-5, 1,2-Benzenediamine, reactions

123-11-5, p-Anisaldehyde, reactions 960-16-7, Tributylphenyltin

6393-40-4, Benzonitrile, 4-amino-3-nitro 17626-40-3, Benzonitrile,

3,4-diamino 17997-47-6, 2-(Tributylstannyl)pyridine 24850-33-7,

Allyltributyltin 59020-10-9, 3-(Tributylstannyl)pyridine 124252-41-1,

4-(Tributylstannyl)pyridine

RL: RCT (Reactant); RACT (Reactant or reagent)

(preparation of terbenzimidazoles as topoisomerase I inhibitors)

IT 875-51-4P, Benzenamine, 4-bromo-2-nitro 4085-18-1P, [1,1'-Biphenyl]-4-

amine, 3-nitro 58442-17-4P, 1H-Benzimidazole-5-carboxaldehyde

59656-62-1P, Benzenamine, 2-nitro-4-(4-pyridinyl) 126824-22-4P

160522-85-0P, Benzenamine, 2-nitro-4-(2-propenyl) 167959-16-2P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation of terbenzimidazoles as topoisomerase I inhibitors)


IT 167959-21-9P, 2,5':2',5''-Ter-1H-benzimidazole

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(2,5':2',5''-Ter-1H-benzimidazole; preparation of terbenzimidazoles as topoisomerase I inhibitors)


RN 167959-21-9 HCAPLUS

CN 2,5':2',5''-Ter-1H-benzimidazole (9CI) (CA INDEX NAME)




=> d all fhitr 127 tot

L27 ANSWER 1 OF 3 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 2003:576172 HCAPLUS
 DN 139:261219
 ED Entered STN: 29 Jul 2003
 TI Influence of Phenyl Ring Disubstitution on Bisbenzimidazole and Terbenzimidazole Cytotoxicity: Synthesis and Biological Evaluation as Radioprotectors
 AU Tawar, Urmila; Jain, Akash K.; Dwarakanath, B. S.; Chandra, Ramesh; Singh, Yogendra; Chaudhury, N. K.; Khaitan, Divya; Tandon, Vibha
 CS Dr. B. R. Ambedkar Center for Biomedical Research, University of Delhi, Delhi, 110007, India
 SO Journal of Medicinal Chemistry (2003), 46(18), 3785-3792
 CODEN: JMCMAR; ISSN: 0022-2623
 PB American Chemical Society
 DT Journal
 LA English
 CC 28-9 (Heterocyclic Compounds (More Than One Hetero Atom))
 Section cross-reference(s): 1
 OS CASREACT 139:261219
 GI



I



II

AB In a search of non-toxic and non-mutagenic DNA radioprotectors, two new disubstituted benzimidazoles I and II were synthesized. The radiomodifying effects of I and II were investigated with a human glioma cell line exposed to low linear energy transfer radiation by determining cell survival and cell proliferation compared with effects of the parent compound, Hoechst 33342. Cytotoxicity assayed by analyzing clonogenicity, cell growth, and metabolic viability showed that both I and II were non-toxic at 100 μ M after 72 h of exposure, whereas Hoechst 33342 resulted in lysis of 77% of these cells in 24 h. Macrocolony assay (clonogenicity) showed that 73%, 92%, and 10% of the cells survived when treated with 100 μ M I, II, and Hoechst 33342, resp. Both I and II did not affect the growth of BMG-1 cells. At 10 μ M, I

Search done by Noble Jarrell

- and II showed 82% and 37% protection against radiation-induced cell death (macrocolony assay) while 100% protection was observed against growth inhibition. Disubstitution of the Ph ring has not only reduced cytotoxicity but also enhanced DNA-ligand stability, conferring high degree of radioprotection.
- ST benzimidazole bis disubstituted prepn cytotoxicity DNA binding radioprotective; terbenzimidazole disubstituted prepn cytotoxicity DNA binding radioprotective
- IT Structure-activity relationship
(DNA-binding; preparation, cytotoxicity, DNA binding properties and radioprotective effects of piperazinyl bisbenzimidazole and terbenzimidazole)
- IT Radiation
(damage; preparation, cytotoxicity, DNA binding properties and radioprotective effects of piperazinyl bisbenzimidazole and terbenzimidazole)
- IT Cytotoxicity
Human
Radioprotectants
(preparation, cytotoxicity, DNA binding properties and radioprotective effects of piperazinyl bisbenzimidazole and terbenzimidazole)
- IT 17626-40-3P, 3,4-Diaminobenzonitrile
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(heterocyclization with aldehyde; preparation, cytotoxicity, DNA binding properties and radioprotective effects of piperazinyl bisbenzimidazole and terbenzimidazole)
- IT 120-14-9, 3,4-Dimethoxybenzaldehyde 121-33-5, 4-Hydroxy-3-methoxybenzaldehyde
RL: RCT (Reactant); RACT (Reactant or reagent)
(heterocyclization with diamine; preparation, cytotoxicity, DNA binding properties and radioprotective effects of piperazinyl bisbenzimidazole and terbenzimidazole)
- IT 188860-26-6P 601473-44-3P
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
(preparation, cytotoxicity, DNA binding properties and radioprotective effects of piperazinyl bisbenzimidazole and terbenzimidazole)
- IT 23491-49-8P 601473-40-9P 601473-43-2P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation, cytotoxicity, DNA binding properties and radioprotective effects of piperazinyl bisbenzimidazole and terbenzimidazole)
- IT 6393-40-4, 4-Amino-3-nitrobenzonitrile 23623-05-4
RL: RCT (Reactant); RACT (Reactant or reagent)
(reduction; preparation, cytotoxicity, DNA binding properties and radioprotective effects of piperazinyl bisbenzimidazole and terbenzimidazole)

RE.CNT 62 THERE ARE 62 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Adhikary, A; Int J Radiat Biol 2000, V76, P1157 HCAPLUS
- (2) Adhikary, A; Radioprotection 1997, V32, P89
- (3) Bathini, Y; Chem Res Toxicol 1990, V3, P268 HCAPLUS
- (4) Beerman, T; Biochim Biophys Acta 1992, V1131, P53 HCAPLUS
- (5) Bostock-Smith, C; Nucleic Acids Res 1998, V26, P1660 HCAPLUS
- (6) Capranico, G; Trends Pharmacol Sci 1991, V18, P323
- (7) Chen, A; Cancer Res 1993, V53, P1332 HCAPLUS
- (8) Chen, A; Proc Natl Acad Sci U S A 1993, V90, P8131 HCAPLUS
- (9) Clark, G; Biochemistry 1996, V35, P13745 HCAPLUS
- (10) Clark, G; Nucleic Acids Res 1997, V25, P510
- (11) Czarny, A; J Am Chem Soc 1995, V117, P4716 HCAPLUS
- (12) de, C; Biochemistry 1989, V28, P7849
- (13) Denison, L; Int J Radiat Biol 1992, V61, P69 HCAPLUS
- (14) Dwarakanath, B; Int J Radiat Oncol Biol Phys 1986, V17, P1033
- (15) Ebrahimi, S; Chem Commun 1991, P385
- (16) Ebrahimi, S; J Chem Soc, Chem Commun 1992, P1398 HCAPLUS
- (17) Embrey, K; FEBS Lett 1993, V211, P437 HCAPLUS
- (18) Embrey, K; J Chem Soc, Chem Commun 1991, P1770 HCAPLUS
- (19) Fede, A; Biochemistry 1991, V30, P11377 HCAPLUS
- (20) Fede, A; Structure 1993, V1, P177 HCAPLUS
- (21) Ganapathi, R; Curr Pharm Des 2002, V8, P1945 HCAPLUS
- (22) Gavathiotis, E; Nucleic Acids Res 2000, V28, P728 HCAPLUS
- (23) Haq, I; J Mol Biol 1997, V271, P244 HCAPLUS
- (24) Hilwig, I; Exp Cell Res 1973, V81, P474 HCAPLUS
- (25) Ji, Y; Bioorg Med Chem 2001, V9, P2905 HCAPLUS
- (26) Kallio, M; Mutagenesis 1993, V8, P561 HCAPLUS

Search done by Noble Jarrell


- (27) Kelly, D; Aust J Chem 1994, V47, P247 HCAPLUS
 (28) Kim, J; Bioorg Med Chem 1998, V6, P163 HCAPLUS
 (29) Kim, J; J Med Chem 1996, V39, P992 HCAPLUS
 (30) Kim, J; J Med Chem 1997, V40, P2818 HCAPLUS
 (31) Kumar, S; J Biomol Struct Dyn 1990, V8, P331 HCAPLUS
 (32) Loontjens, F; Biochemistry 1990, V29, P9029 HCAPLUS
 (33) Loontjens, F; Biochemistry 1991, V30, P182 HCAPLUS
 (34) Lyubimova, N; Br J Radiol 2001, V74, P77 HCAPLUS
 (35) Martin, R; Int J Radiat Oncol Biol Phys 1992, V23, P579 HCAPLUS
 (36) Martin, R; Int J Radiat Oncol Biol Phys 1998, V42, P827 HCAPLUS
 (37) Mekapati, S; Bioorg Med Chem 2001, V9, P2885 HCAPLUS
 (38) Mosmann, T; J Immunol Methods 1983, V65, P55 MEDLINE
 (39) Parkinson, J; Biochemistry 1990, V29, P10181 HCAPLUS
 (40) Parkinson, J; Biochemistry 1994, V33, P8442 HCAPLUS
 (41) Pilch, D; Proc Natl Acad Sci U S A 1997, V94, P13565 HCAPLUS
 (42) Pjura, P; J Mol Biol 1987, V197, P257 HCAPLUS
 (43) Portugal, J; Biochim Biophys Acta 1987, V949, P158
 (44) Quintana, J; Biochemistry 1991, V30, P10294 HCAPLUS
 (45) Searle, M; Nucleic Acids Res 1990, V18, P3753 HCAPLUS
 (46) Singh, A; Anti-Cancer Drug Des 2000, V15, P265 HCAPLUS
 (47) Singh, M; Chem Res Toxicol 1992, V5, P597 HCAPLUS
 (48) Singh, S; Indian J Exp Biol 1998, V36, P375 HCAPLUS
 (49) Sorenson, J; Curr Med Chem 2002, V9, P639 HCAPLUS
 (50) Sriram, M; EMBO J 1992, V11, P225 HCAPLUS
 (51) Stokke, T; J Histochem Cytochem 1985, V33, P333 HCAPLUS
 (52) Sun, Q; Bioorg Med Chem Lett 1994, V4, P2871 HCAPLUS
 (53) Sun, Q; J Med Chem 1995, V38, P3638 HCAPLUS
 (54) Tawar, U; Submitted for publication 2002
 (55) Turner, P; Mutat Res 1996, V355, P141 HCAPLUS
 (56) Vega, M; Eur J Biochem 1994, V222, P721 HCAPLUS
 (57) Wang, E; Biochim Biophys Acta 2000, V1481, P63 HCAPLUS
 (58) Wang, H; Anti-Cancer Drug Des 1994, P145
 (59) Wood, A; Nucleic Acids Res 1995, V23, P3678 HCAPLUS
 (60) Xu, Z; Biochemistry 1998, V37, P3558 HCAPLUS
 (61) Yadagiri, B; Synth Commun 1990, V20, P955 HCAPLUS
 (62) Young, S; Br J Cancer 1989, V60, P715 HCAPLUS

IT 601473-44-3P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (preparation, cytotoxicity, DNA binding properties and radioprotective effects of piperazinyl bisbenzimidazole and terbenzimidazole)

RN 601473-44-3 HCAPLUS

CN Phenol, 2-methoxy-4-[5-(4-methyl-1-piperazinyl)[2,5':2',5''-ter-1H-benzimidazol]-2''-yl]- (9CI) (CA INDEX NAME)



L27 ANSWER 2 OF 3 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 2003:463129 HCAPLUS

DN 140:174411

ED Entered STN: 17 Jun 2003

TI Some new bi- and ter-benzimidazole derivatives as topoisomerase I inhibitors

AU Alper, Sabiha; Temiz Arpacı, Ozlem; Aki-Sener, Esin; Yalcin, Ismail
 CS Faculty of Pharmacy, Department of Pharmaceutical Chemistry, Ankara University Tandoğan, Ankara, 06100, Turk.

SO Farmaco (2003), 58(7), 497-507

CODEN: FRMCE8; ISSN: 0014-827X

PB Editions Scientifiques et Medicales Elsevier

DT Journal

LA English

CC 1-3 (Pharmacology)

AB The discovery of DNA topoisomerases has added a new dimension to the study of anticancer drugs. In the last years detailed investigation

of bi- and ter-benzimidazole derivs. revealed that these compds. are a new class of topoisomerase I inhibitors that poisons mammalian topoisomerase I. In this context a survey about topoisomerase I poisoning activity and cytotoxicity of bi- and ter-benzimidazoles is given. Moreover some recent results about new derivs., some structure-activity relationships and comparison of activity of various functional groups are discussed.

ST topoisomerase I inhibitor benzimidazole deriv

IT Antitumor agents

Structure-activity relationship

(some new bi- and ter-benzimidazole derivs. as topoisomerase I inhibitors and antitumor activity)

IT 143180-75-0

RL: BSU (Biological study, unclassified); BIOL (Biological study)

(some new bi- and ter-benzimidazole derivs. as topoisomerase I inhibitors and antitumor activity)

IT 2620-81-7 23491-54-5 23491-55-6 23554-98-5 23623-08-7 96954-35-7

96954-36-8 126824-21-3 126824-22-4 167959-13-9 167959-14-0

167959-15-1 167959-17-3 167959-21-9,

2,5':2',5''-Ter-1H-benzimidazole 167959-22-0 167959-23-1

167959-24-2 167959-25-3 167959-26-4

167959-27-5 174422-17-4 174648-30-7 174648-31-8

174648-32-9 174648-33-0 174648-34-1 174648-35-2 174648-36-3

174648-37-4 174648-38-5 174648-39-6 174648-40-9 174648-41-0

174648-42-1 174648-43-2 178970-15-5 178970-16-6 178970-30-4

185199-36-4 185199-38-6 192879-67-7

192879-68-8 205749-93-5 205749-94-6

205749-95-7 205749-96-8 205749-97-9,

[2,5':2',5''-Ter-1H-benzimidazol]-5-ol 205749-98-0

230308-96-0 230308-97-1 230308-98-2 237429-52-6

237429-53-7 237429-54-8 237429-55-9

237429-56-0 237429-57-1 237429-58-2

237429-59-3 277754-98-0 277754-99-1

319916-61-5

RL: PAC (Pharmacological activity); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(some new bi- and ter-benzimidazole derivs. as topoisomerase I inhibitors and antitumor activity)

RE.CNT 58 THERE ARE 58 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

(1) Allan, Y; Annu Rev Pharmacol Toxicol 1994, V34, P191

(2) Anizon, F; J Med Chem 1997, V40, P3456 HCAPLUS

(3) Anon; Molecular biology of DNA Topoisomerases and its Application to Chemotherapy 1993, P377

(4) Capranico, G; TIPS 1997, V18, P323 HCAPLUS

(5) Champoux, J; Proc Natl Acad Sci USA 1972, V69, P143 HCAPLUS

(6) Chen, A; Cancer Res 1993, V53, P1332 HCAPLUS

(7) Chen, A; Proc Natl Acad Sci USA 1993, V90, P8131 HCAPLUS

(8) Finlay, J; Eur J Cancer 1990, V26, P586

(9) Fujii, N; J Biol Chem 1993, V268, P13160 HCAPLUS

(10) Gellert, M; Proc Natl Acad Sci USA 1976, V73, P3872 HCAPLUS

(11) Hertzberg, R; Biochemistry 1989, V28, P4629 HCAPLUS

(12) Hertzberg, R; J Biol Chem 1990, V265, P19287 HCAPLUS

(13) Hertzberg, R; J Med Chem 1989, V32, P715 HCAPLUS

(14) Hsiang, Y; J Biol Chem 1985, V260, P14873 HCAPLUS

(15) Jin, S; Bioorg Med Chem Lett 2000, V10, P719 HCAPLUS

(16) Kim, J; Bioorg Med Chem 1996, V4, P621 HCAPLUS

(17) Kim, J; Bioorg Med Chem 1998, V6, P163 HCAPLUS

(18) Kim, J; J Med Chem 1996, V39, P992 HCAPLUS

(19) Kim, J; J Med Chem 1997, V40, P2818 HCAPLUS

(20) Kuhholzer, B; Anim Reprod Sci 2001, V66, P109 HCAPLUS

(21) Liu, L; Annu Rev Biochem 1989, V58, P351 HCAPLUS

(22) Liu, L; Important Adv Oncol 1992, V1992, P79

(23) Liu, L; Nature 1979, V281, P456 HCAPLUS

(24) McHugh, M; Biochem Pharmacol 1989, V38, P2323 HCAPLUS

(25) Nelson, E; Proc Natl Acad Sci USA 1984, V81, P1361 HCAPLUS

(26) Nitiss, J; Biochem Biophys Acta 1998, V1400, P63 HCAPLUS

(27) Oren, I; Arzneimittel-Forschung/Drug Res 1997, V47, P1393 MEDLINE

(28) Oren, I; Eur J Pharm Sci 1998, V7, P153

(29) Pindur, U; Pharmazie 1998, V53, P79 HCAPLUS

(30) Poddevin, B; Mol Pharmacol 1993, V44, P767 HCAPLUS

(31) Pommier, Y; Proc Natl Acad Sci USA 1995, V92, P8861 HCAPLUS

(32) Porter, S; Nucleic Acids Res 1989, V17, P8521 HCAPLUS

(33) Rangarajan, M; Bioorg Med Chem 2000, V8, P1371 HCAPLUS

(34) Rangarajan, M; Bioorg Med Chem 2000, V8, P2591 HCAPLUS

(35) Riou, J; Cancer Res 1993, V53, P5987 HCAPLUS

(36) Riou, J; Mol Pharmacol 1991, V40, P699 HCAPLUS

Search done by Noble Jarrell


- (37) Rowe, T; Cancer Res 1986, V46, P2021 HCAPLUS
- (38) Settimo, A; Eur J Med Chem 1998, V33, P685
- (39) Skibo, E; J Med Chem 1997, V40, P1327 HCAPLUS
- (40) Skibo, E; J Med Chem 2000, V43, P629
- (41) Sun, Q; Bioorg Med Chem Lett 1994, V4, P2871 HCAPLUS
- (42) Sun, Q; J Med Chem 1995, V38, P3638 HCAPLUS
- (43) Temiz, O; Il Farmaco 1998, V53, P337 HCAPLUS
- (44) Tewey, K; Science 1984, V226, P466 HCAPLUS
- (45) Tewey, K; J Biol Chem 1984, V259, P9182
- (46) Tobey, R; Proc Natl Acad Sci USA 1990, V87, P5104 HCAPLUS
- (47) Tolner, B; Mol Pharmacol 2001, V59, P699 HCAPLUS
- (48) Trask, D; Proc Natl Acad Sci USA 1988, V85, P1417 HCAPLUS
- (49) Wall, M; J Am Chem Soc 1966, V88, P3888 HCAPLUS
- (50) Wang, J; J Mol Biol 1971, V55, P523 HCAPLUS
- (51) Wassermann, K; Mol Pharmacol 1990, V38, P38 HCAPLUS
- (52) Woynarowski, J; Mol Pharmacol 1989, V35, P177 HCAPLUS
- (53) Xu, Z; Biochemistry 1998, V37, P3558 HCAPLUS
- (54) Yalcin, I; Eur J Med Chem 1990, V25, P705 HCAPLUS
- (55) Yamashita, Y; Biochemistry 1992, V31, P12069 HCAPLUS
- (56) Yoshinari, T; Cancer Res 1993, V53, P490 HCAPLUS
- (57) Zhang, X; Ann Clin Lab Sci 2001, V31, P187 HCAPLUS
- (58) Zhou, R; J Med Chem 1996, V39, P4321 HCAPLUS

IT 167959-17-3

RL: PAC (Pharmacological activity); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(some new bi- and ter-benzimidazole derivs. as topoisomerase I inhibitors and antitumor activity)

RN 167959-17-3 HCAPLUS

CN [2,5':2',5''-Ter-1H-benzimidazole]-5-carbonitrile, 2''-(4-methoxyphenyl)-(9CI) (CA INDEX NAME)



L27 ANSWER 3 OF 3 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 2001:845775 HCAPLUS

DN 136:128592

ED Entered STN: 21 Nov 2001

TI Comparative QSAR studies on bisbenzimidazoles and terbenzimidazoles inhibiting topoisomerase I

AU Mekapati, Suresh Babu; Hansch, Corwin

CS Department of Chemistry, Pomona College, Claremont, CA, 91711, USA

SO Bioorganic & Medicinal Chemistry (2001), 9(11), 2885-2893

CODEN: BMECEP; ISSN: 0968-0896

PB Elsevier Science Ltd.

DT Journal

LA English

CC 1-3 (Pharmacology)

AB Terbenzimidazoles that inhibit topoisomerase are of interest as anticancer drugs. We have reviewed the literature and have developed 13 quant. structure-activity relationships (QSARs) on cleaving DNA or inhibiting the growth of tumor cell cultures. The results are correlated with octanol/water partition coeffs. or mol. refractivity. Suggestions have been made for the development of improved derivs.

ST antitumor bisbenzimidazole terbenzimidazole QSAR topoisomerase I; lymphoblastoma bisbenzimidazole terbenzimidazole QSAR topoisomerase I

IT DNA

RL: BSU (Biological study, unclassified); BIOL (Biological study)
(cleavage; comparative QSAR studies on bisbenzimidazoles and terbenzimidazoles inhibiting topoisomerase I)

IT Antitumor agents

Human

Lymphoma

QSAR (structure-activity relationship)

Search done by Noble Jarrell

(comparative QSAR studies on bisbenzimidazoles and terbenzimidazoles inhibiting topoisomerase I)

IT 143180-75-0
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (comparative QSAR studies on bisbenzimidazoles and terbenzimidazoles inhibiting topoisomerase I)

IT 2620-81-7 23491-52-3 23491-54-5 23491-55-6 23554-98-5 23623-06-5
 23623-08-7 54998-13-9 96954-35-7 96954-36-8 126824-21-3
 126824-22-4 160522-67-8 160522-68-9 160522-69-0 160522-70-3
 160522-71-4 160522-72-5 167959-13-9 167959-14-0 167959-15-1
 167959-21-9, 2,5':2',5''-Ter-1H-benzimidazole 167959-22-0
 167959-23-1 167959-24-2 167959-25-3
 167959-26-4 167959-27-5 174648-31-8 174648-32-9
 174648-33-0 174648-34-1 174648-35-2 174648-36-3 174648-37-4
 174648-38-5 174648-39-6 174648-40-9 174648-41-0 174648-42-1
 174648-43-2 178970-15-5 178970-16-6 178970-30-4 185199-36-4
 185199-38-6 185199-39-7 192879-62-2
 192879-64-4 192879-67-7 192879-68-8
 192879-69-9 192879-73-5 192879-74-6
 192879-75-7 192879-76-8 192879-77-9
 192879-78-0 205749-93-5 205749-94-6
 205749-95-7 205749-96-8 205749-97-9,
 [2,5':2',5''-Ter-1H-benzimidazol]-5-ol 205749-98-0
 237429-45-7 237429-46-8 237429-48-0
 237429-52-6 237429-53-7 237429-54-8
 237429-55-9 237429-56-0 237429-57-1
 237429-58-2 237429-59-3 288579-81-7
 319916-61-5 392287-14-8 392287-15-9
 392287-16-0 392287-17-1 392287-18-2
 392287-19-3 392287-20-6 392287-21-7
 392287-22-8 392287-23-9
 RL: PAC (Pharmacological activity); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (comparative QSAR studies on bisbenzimidazoles and terbenzimidazoles inhibiting topoisomerase I)

RE.CNT 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD


RE

- (1) Abraham, M; Chromatographica 1987, V23, P243 HCAPLUS
- (2) Chen, A; Annu Rev Pharmacol Toxicol 1994, V34, P191 HCAPLUS
- (3) Chen, A; Cancer Res 1993, V53, P1332 HCAPLUS
- (4) Chen, A; Proc Natl Acad Sci 1993, V90, P8131 HCAPLUS
- (5) D'Arpa, P; Biochem Biophys Acta 1989, V989, P163 HCAPLUS
- (6) Gao, H; Chem Rev 1999, V99, P723 HCAPLUS
- (7) Gao, H; Drug Metab Rev 1996, V18, P513
- (8) Garg, R; Chem Rev 1999, V99, P3525 HCAPLUS
- (9) Hansch, C; Acc Chem Res 1969, V2, P232 HCAPLUS
- (10) Hansch, C; Chem Rev 1996, V96, P1045 HCAPLUS
- (11) Hansch, C; Chem Rev 1997, V97, P2995 HCAPLUS
- (12) Hansch, C; Exploring QSAR Hydrophobic, Electronic and Steric Constants 1995
- (13) Hansch, C; J Pharm Sci 1987, V76, P663 HCAPLUS
- (14) Hansch, C; Nature 1962, V194, P178 HCAPLUS
- (15) Hansch, C; Quant Struct-Act Relat 1990, V9, P223 HCAPLUS
- (16) Kim, J; Bioorg Med Chem 1996, V4, P621 HCAPLUS
- (17) Kim, J; Bioorg Med Chem 1998, V6, P163 HCAPLUS
- (18) Kim, J; J Med Chem 1996, V39, P992 HCAPLUS
- (19) Kim, J; J Med Chem 1997, V40, P2818 HCAPLUS
- (20) Koo, H; Proc Natl Acad Sci 1991, V88, P1212 HCAPLUS
- (21) Kurup, A; Chem Rev 2000, V100, P909 HCAPLUS
- (22) Liu, L; Annu Rev Biochem 1989, V58, P351 HCAPLUS
- (23) Liu, L; Proc Natl Acad Sci 1987, V84, P7024 HCAPLUS
- (24) Rangarajan, M; Bioorg Med Chem 2000, V8, P1371 HCAPLUS
- (25) Rangarajan, M; Bioorg Med Chem 2000, V8, P2591 HCAPLUS
- (26) Schneider, E; Adv Pharmacol 1990, V21, P149 HCAPLUS
- (27) Sun, Q; Bioorg Med Chem Lett 1994, V4, P2871 HCAPLUS
- (28) Sun, Q; J Med Chem 1995, V38, P3638 HCAPLUS
- (29) Tsao, Y; Cell 1989, V56, P111 HCAPLUS
- (30) Wang, J; Annu Rev Biochem 1985, V54, P665 MEDLINE

IT 167959-21-9, 2,5':2',5''-Ter-1H-benzimidazole
 RL: PAC (Pharmacological activity); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (comparative QSAR studies on bisbenzimidazoles and terbenzimidazoles inhibiting topoisomerase I)

RN 167959-21-9 HCAPLUS

CN 2,5':2',5''-Ter-1H-benzimidazole (9CI) (CA INDEX NAME)



=> d all hitstr 128 tot

L28 ANSWER 1 OF 17 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 2004:606451 HCAPLUS
 DN 141:157115
 ED Entered STN: 29 Jul 2004
 TI A process for the synthesis of bisbenzimidazole derivatives, useful as
 radioprotective agents
 IN Jain, Akash; Tawar, Urmila; Chandra, Ramesh; Dwarakanath, B. s.;
 Chaudhury, N. K.
 PA University of Delhi, India; Tandon, Vibha
 SO PCT Int. Appl., 25 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 IC ICM C07D235-18
 ICS C07D235-20
 CC 28-9 (Heterocyclic Compounds (More Than One Hetero Atom))
 Section cross-reference(s): 1, 8
 FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004063170	A1	20040729	WO 2003-IN301	20030908
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
PRAI IN 2003-DE32	A	20030109		

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
WO 2004063170	ICM	C07D235-18
	ICS	C07D235-20


OS CASREACT 141:157115
 GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The invention relates to a preparation of bisbenzimidazole derivs., e.g. I,
 useful as radioprotective agents. The invented compds. are DNA binding
 ligands (Minor Groove Binding Ligands) that afford radioprotective effect
 without significant mutagenicity and cytotoxic effects. Cell survival
 assay showed that I has 73% cell survival at 100 .mu.M concentration For
 instance, I was prepared via heterocyclization of 3,4-dimethoxybenzaldehyde
 and benzimidazole derivative II with a yield of 30% (example 10).
 ST bisbenzimidazole prepn radioprotectant radiation; dimethoxybenzaldehyde
 diamine heterocyclization
 IT Heterocyclization
 Human
 Radioprotectants
 (process for the synthesis of bisbenzimidazole derivs., useful as
 radioprotective agents)
 IT Radiation
 (treatment of; process for the synthesis of bisbenzimidazole derivs.,
 useful as radioprotective agents)

Search done by Noble Jarrell

- IT 188860-26-6P 601473-44-3P
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (process for the synthesis of bisbenzimidazole derivs., useful as radioprotective agents)
- IT 109-01-3, 1-Methylpiperazine 121-33-5, 3-Methoxy-4-hydroxybenzaldehyde 588-07-8, m-Chloroacetanilide 5443-33-4 17626-40-3, 3,4-Diaminobenzonitrile 29289-18-7
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (process for the synthesis of bisbenzimidazole derivs., useful as radioprotective agents)
- IT 120-14-9P, 3,4-Dimethoxybenzaldehyde 1635-61-6P, 2-Nitro-5-chloroaniline 6393-40-4P 23491-48-7P 23491-49-8P 23623-05-4P 54998-08-2P 54998-39-9P 165596-29-2P 601473-40-9P 601473-43-2P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (process for the synthesis of bisbenzimidazole derivs., useful as radioprotective agents)
- IT 601473-44-3P
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (process for the synthesis of bisbenzimidazole derivs., useful as radioprotective agents)
- RN 601473-44-3 HCAPLUS
- CN Phenol, 2-methoxy-4-[5-(4-methyl-1-piperazinyl)[2,5':2',5''-ter-1H-benzimidazol]-2''-yl]- (9CI) (CA INDEX NAME)



- L28 ANSWER 2 OF 17 HCAPLUS COPYRIGHT 2005 ACS on STN
- AN 2003:877186 HCAPLUS
- DN 140:93976
- ED Entered STN: 10 Nov 2003
- TI Synthesis of a terbenzimidazole topoisomerase I poison via iterative borinate ester couplings
- AU Wang, Ben B.; Smith, Paul J.
- CS Department of Chemistry and Biochemistry, University of Maryland, Baltimore, MD, 21250, USA
- SO Tetrahedron Letters (2003), 44(50), 8967-8969
 CODEN: TELEAY; ISSN: 0040-4039
- PB Elsevier Science B.V.
- DT Journal
- LA English
- CC 28-9 (Heterocyclic Compounds (More Than One Hetero Atom))
- AB A concise, efficient synthesis is described for a terbenzimidazole that acts as a potent topoisomerase I poison. The strategy involves iterative palladium-catalyzed borinate ester cross-couplings and should be applicable to the synthesis of analogs containing heterocycles other than benzimidazole.
- ST terbenzimidazole topoisomerase poison coupling reaction borinate ester palladium catalyst
- IT Cross-coupling reaction
 Cross-coupling reaction catalysts
 (preparation of terbenzimidazole topoisomerase I poison via iterative palladium-catalyzed cross-coupling reaction of corresponding borinate ester)
- IT 14221-01-3, Tetrakis(triphenylphosphine)palladium 72287-26-4, Dichloro(diphenylphosphinoferrocene)palladium
 RL: CAT (Catalyst use); USES (Uses)
 (preparation of terbenzimidazole topoisomerase I poison via iterative palladium-catalyzed cross-coupling reaction of corresponding borinate ester)

IT 98-80-6, Phenylboronic acid 21304-38-1 73183-34-3
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (preparation of terbenzimidazole topoisomerase I poison via iterative
 palladium-catalyzed cross-coupling reaction of corresponding borinate
 ester)

IT 78597-27-0P 645414-73-9P 645414-74-0P 645414-75-1P 645414-76-2P
 645414-77-3P 645414-78-4P 645414-79-5P 645414-80-8P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
 (Reactant or reagent)
 (preparation of terbenzimidazole topoisomerase I poison via iterative
 palladium-catalyzed cross-coupling reaction of corresponding borinate
 ester)

IT 167959-27-5P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of terbenzimidazole topoisomerase I poison via iterative
 palladium-catalyzed cross-coupling reaction of corresponding borinate
 ester)

RE.CNT 19 THERE ARE 19 CITED REFERENCES AVAILABLE FOR THIS RECORD


RE

- (1) Beerman, T; Biochim Biophys Res Acta 1992, V1131, P53 HCAPLUS
- (2) Bellamy, F; Tetrahedron Lett 1984, V25, P839 HCAPLUS
- (3) Bergman, J; J Org Chem 1992, V57, P2495 HCAPLUS
- (4) Chen, A; Proc Natl Acad Sci USA 1993, V90, P8131 HCAPLUS
- (5) Ezquerro, J; Tetrahedron 1997, V53, P2755
- (6) Gros, P; Org Lett 2002, V4, P1759 HCAPLUS
- (7) Ishiyama, T; J Org Chem 1995, V60, P7508 HCAPLUS
- (8) Jaxel, C; J Biol Chem 1991, V266, P20418 HCAPLUS
- (9) Kissinger, K; Biochemistry 1987, V26, P5590 HCAPLUS
- (10) Leonard, N; J Org Chem 1979, V44, P4612 HCAPLUS
- (11) Lown, J; Advances in DNA Sequence-Specific Agents 1998, V3, P67 HCAPLUS
- (12) Nicolaou, K; Nature 1997, V387, P268 HCAPLUS
- (13) Pieltre, S; Tetrahedron Lett 1997, V38, P1197 HCAPLUS
- (14) Pjura, P; J Mol Biol 1987, V197, P257 HCAPLUS
- (15) Singh, M; Chem Res Toxicol 1992, V5, P597 HCAPLUS
- (16) Sun, Q; J Med Chem 1995, V38, P3638 HCAPLUS
- (17) Villemin, D; Synth Commun 1996, V26, P2895 HCAPLUS
- (18) Wilson, J; Aust J Chem 1983, V36, P2317 HCAPLUS
- (19) Xu, Z; Biochemistry 1998, V37, P3558 HCAPLUS

IT 645414-80-8P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
 (Reactant or reagent)
 (preparation of terbenzimidazole topoisomerase I poison via iterative
 palladium-catalyzed cross-coupling reaction of corresponding borinate
 ester)

RN 645414-80-8 HCAPLUS


CN 2,5':2',5''-Ter-1H-benzimidazole, 6-phenyl-1,1',1''-tris(phenylmethyl)-
 (9CI) (CA INDEX NAME)



IT 167959-27-5P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of terbenzimidazole topoisomerase I poison via iterative
 palladium-catalyzed cross-coupling reaction of corresponding borinate
 ester)

RN 167959-27-5 HCAPLUS

CN 2,5':2',5''-Ter-1H-benzimidazole, 5-phenyl- (9CI) (CA INDEX NAME)



L28 ANSWER 3 OF 17 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 2003:826193 HCAPLUS
 DN 139:376009
 ED Entered STN: 22 Oct 2003
 TI Minor Groove Binding DNA Ligands with Expanded A/T Sequence Length
 Recognition, Selective Binding to Bent DNA Regions and Enhanced
 Fluorescent Properties
 AU Tawar, Urmila; Jain, Akash K.; Chandra, Ramesh; Singh, Yogendra;
 Dwarakanath, B. S.; Chaudhury, N. K.; Good, Liam; Tandon, Vibha
 CS Dr. B. R. Ambedkar Center for Biomedical Research, University of Delhi,
 Delhi, 110007, India
 SO Biochemistry (2003), 42(45), 13339-13346
 CODEN: BICHAW; ISSN: 0006-2960
 PB American Chemical Society
 DT Journal
 LA English
 CC 3-3 (Biochemical Genetics)
 AB DNA minor groove ligands provide a paradigm for double-stranded DNA
 recognition, where common structural motifs provide a crescent shape that
 matches the helix turn. Since minor groove ligands are useful in
 medicine, new ligands with improved binding properties based on the
 structural information about DNA-ligand complexes could be useful in
 developing new drugs. Here, two new synthetic analogs of AT specific
 Hoechst 33258 5-(4-methylpiperazin-1-yl)-2-[2'-(3,4-dimethoxyphenyl)-5'-
 benzimidazolyl] benzimidazole (DMA) and 5-(4-methylpiperazin-1-yl)-2-
 [2'-(4-hydroxy-3-methoxyphenyl)-5'-benzimidazolyl]-5'-benzimidazolyl]
 benzimidazole (TBZ) were evaluated for their DNA binding properties. Both
 analogs are substituted on the Ph ring. DMA contains two ortho positioned
 methoxy groups, and TBZ contains a phenolic group at C-4 and a methoxy
 group at C-3. Fluorescence yield upon DNA binding increased 100-fold for
 TBZ and 16-fold for DMA. Like the parent compound, the new ligands showed
 low affinity to GC-rich (K .apprx. 4 .times. 10⁷ M⁻¹) relative to
 AT-rich sequences (K .apprx. 5 .times. 10⁸ M⁻¹), and fluorescence
 lifetime and anisotropy studies suggest two distinct DNA-ligand complexes.
 Binding studies indicate expanded sequence recognition for TBZ (8-10 AT
 base pairs) and tighter binding (.DELTA.Tm of 23 .degree.C for d
 (GA5T5C)). Finally, EMSA and equilibrium binding titration studies indicate that
 TBZ preferentially binds highly hydrated duplex domains with altered
 A-tract conformations d (GA4T4C)₂ (K = 3.55 .times. 10⁹ M⁻¹) and alters
 its structure over d (GT4A4C)₂ (K = 3.3 .times. 10⁸ M⁻¹) sequences.
 Altered DNA structure and higher fluorescence output for the bound
 fluorophore are consistent with adaptive binding and a constrained final
 complex. Therefore, the new ligands provide increased sequence and
 structure selective recognition and enhanced fluorescence upon minor
 groove binding, features that can be useful for further development as
 probes for chromatin structure stability.
 ST DNA ligand AT minor groove helix conformation fluorescence
 IT Genetic element
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (AT-rich element; minor groove-binding DNA ligands recognizes AT-rich
 sequence and enhances fluorescence)
 IT Conformation
 Helix (conformation)
 (DNA; minor groove-binding DNA ligands recognizes AT-rich sequence and
 enhances fluorescence)
 IT Fluorescence
 (minor groove-binding DNA ligands recognizes AT-rich sequence and
 enhances fluorescence)
 IT DNA
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (minor groove-binding DNA ligands recognizes AT-rich sequence and
 enhances fluorescence)
 IT 188860-26-6 601473-44-3
 RL: BSU (Biological study, unclassified); BUU (Biological use,
 unclassified); BIOL (Biological study); USES (Uses)
 (minor groove-binding DNA ligands recognizes AT-rich sequence and
 enhances fluorescence)
 RE.CNT 34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD
 RE
 (1) Arthanari, H; Nucleic Acids Res 1998, V26, P3724 HCAPLUS
 (2) Balagurumorthy, P; J Biol Chem 1994, V269(34), P21858 HCAPLUS
 (3) Benesi, H; J Am Chem Soc 1949, V71, P2703 HCAPLUS
 (4) Bhattacharya, S; Biochim Biophys Acta 1997, V1323, P29 HCAPLUS
 (5) Breusegem, S; J Mol Biol 2001, V308, P649 HCAPLUS


- (6) Burkhoff, A; Nature 1988, V331, P455 MEDLINE
- (7) Caneva, R; Biochim Biophys Acta 1997, V1353, P93 HCAPLUS
- (8) Diekmann, S; J Mol Biol 1985, V186, P1 HCAPLUS
- (9) Eriksson, S; Biochemistry 1993, V32, P2987 HCAPLUS
- (10) Gavathiotis, E; Nucleic Acids Res 2000, V28, P728 HCAPLUS
- (11) Hagerman, P; Annu Rev Biochem 1990, V59, P755 HCAPLUS
- (12) Hertzberg, R; Biochemistry 1984, V23, P3934 HCAPLUS
- (13) Ji, Y; Bioorg Med Chem 2001, V9, P2905 HCAPLUS
- (14) Koo, H; Proc Natl Acad Sci U S A 1988, V85, P1763 HCAPLUS
- (15) Kubista, M; Biochemistry 1987, V26, P4545 HCAPLUS
- (16) Kumar, C; J Am Chem Soc 1993, V115, P8547 HCAPLUS
- (17) Kumar, C; J Chem Soc, Chem Commun 1992, P470 HCAPLUS
- (18) Martin, R; Aust J Chem 1986, V39, P373 HCAPLUS
- (19) Meehan, T; J Biol Chem 1982, V257, P10479 HCAPLUS
- (20) Mekapati, S; Bioorg Med Chem 2001, V9, P2885 HCAPLUS
- (21) Nagaich, A; J Biol Chem 1994, V269, P7824 HCAPLUS
- (22) Pierce, M; Biochemistry 1993, V32, P127
- (23) Pilch, D; Proc Natl Acad Sci U S A 1997, V94, P13565 HCAPLUS
- (24) Sadat, E; Chem Commun 1997, V4, P385
- (25) Schemechel, D; Biopolymers 1971, V10, P465
- (26) Sprou, D; Nucleic Acids Res 1995, V23, P1816 HCAPLUS
- (27) Tabernaro, L; Biochemistry 1993, V32, P8403
- (28) Tawar, U; J Med Chem 2003, V46, P3785 HCAPLUS
- (29) Teng, M; Nucleic Acids Res 1988, V16, P2671 HCAPLUS
- (30) Travers, A; Annu Rev Biochem 1989, V58, P427 HCAPLUS
- (31) Turner, P; Mutation Res 1996, V355, P141 HCAPLUS
- (32) Vega, M; Eur J Biochem 1994, V222, P721 HCAPLUS
- (33) Wang, H; Anticancer Drug Des 1994, V9, P145
- (34) Wu, H; Nature 1984, V308(5), P509

IT 601473-44-3

RL: BSU (Biological study, unclassified); BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
 (minor groove-binding DNA ligands recognizes AT-rich sequence and enhances fluorescence)

RN 601473-44-3 HCAPLUS

CN Phenol, 2-methoxy-4-[5-(4-methyl-1-piperazinyl)[2,5':2',5''-ter-1H-benzimidazol]-2''-yl]- (9CI) (CA INDEX NAME)



L28 ANSWER 4 OF 17 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 2001:845778 HCAPLUS

DN 136:131135

ED Entered STN: 21 Nov 2001

TI Tris-benzimidazole derivatives: design, synthesis and DNA sequence recognition

AU Ji, Yu-Hua; Bur, Daniel; Hasler, Walter; Schmitt, Valerie Runtz; Dorn, Arnulf; Bailly, Christian; Waring, Michael J.; Hochstrasser, Remo; Leupin, Werner

CS Pharma Research Preclinical Gene Technologies and Infectious Diseases, F. Hoffmann-La Roche Ltd, Basel, CH-4070, Switz.

SO Bioorganic & Medicinal Chemistry (2001), 9(11), 2905-2919
 CODEN: BMECEP; ISSN: 0968-0896

PB Elsevier Science Ltd.

DT Journal

LA English

CC 9-14 (Biochemical Methods)

AB Two tris-benzimidazole derivs. have been designed and synthesized based on the known structures of the bis-benzimidazole stain Hoechst 33258 complexed to short oligonucleotide duplexes derived from single crystal x-ray studies and from NMR. In both derivs. the phenol group has been replaced by a methoxy-Ph substituent. Whereas one tris-benzimidazole carries a N-methyl-piperazine at the 6-position, the other one has this group replaced by a 2-amino-pyrrolidine ring. This latter substituent

results in stronger DNA binding. The optimized synthesis of the drugs is described. The two tris-benzimidazoles exhibit high AT-base pair (bp) selectivity evident in footprinting expts. which show that five to six base pairs are protected by the tris-benzimidazoles as compared to four to five protected by the bis-benzimidazoles. The tris-benzimidazoles bind well to sequences like 5'-TAAAC, 5'-TTTAC and 5'-TTTAT, but it is also evident that they can bind weakly to sequences such as 5'-TATGTT-3' where the continuity of an AT stretch is interrupted by a single G.cntdot.C base pair.

ST benzimidazole deriv prepn DNA sequence recognition; DNase footprinting
benzimidazole deriv prepn

IT DNA sequences

(design, synthesis and DNA sequence recognition using
tris-benzimidazole derivs.)

IT 98-95-3, Nitrobenzene, reactions 109-01-3, N-Methylpiperazine
123-11-5, 4-Methoxybenzaldehyde, reactions 528-45-0, 3,4-Dinitrobenzoic
acid 5344-44-5, 5-Chloro-3-nitroaniline 16645-06-0,
Dimethylhydroxylamine hydrochloride 37466-90-3 99724-19-3,
3-tert-Butoxycarbonylaminopyrrolidine

RL: RCT (Reactant); RACT (Reactant or reagent)

(design, synthesis and DNA sequence recognition using
tris-benzimidazole derivs.)

IT 24376-18-9P 126824-19-9P 126824-21-3P 126824-22-4P 167959-16-2P
391903-19-8P 391903-20-1P 391903-21-2P 391903-22-3P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
(Reactant or reagent)

(design, synthesis and DNA sequence recognition using
tris-benzimidazole derivs.)

IT 23491-48-7P 182496-21-5P 391903-23-4P 391903-24-5P
391903-25-6P 391903-26-7P

RL: SPN (Synthetic preparation); PREP (Preparation)

(design, synthesis and DNA sequence recognition using
tris-benzimidazole derivs.)

IT 9003-98-9, DNase

RL: BSU (Biological study, unclassified); BIOL (Biological study)
(footprinting; design, synthesis and DNA sequence recognition using
tris-benzimidazole derivs.)

RE.CNT 54 THERE ARE 54 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Aymami, J; Nucleic Acids Res 1999, V27, P2691 HCAPLUS
- (2) Bailly, C; Bioconjugate Chem 1998, V9, P513 HCAPLUS
- (3) Bailly, C; J Biomol Struct Dyn 1995, V12, P869 HCAPLUS
- (4) Bathini, Y; Chem Res Toxicol 1990, V3, P268 HCAPLUS
- (5) Bathini, Y; Synthetic Commun 1990, V20, P955
- (6) Bostock-Smith, C; Nucleic Acids Res 1998, V26, P1660 HCAPLUS
- (7) Carrondo, M; Biochemistry 1989, V28, P7849 HCAPLUS
- (8) Clark, G; Biochemistry 1996, V35, P13745 HCAPLUS
- (9) Clark, G; Nucleic Acids Res 1996, V24, P4882 HCAPLUS
- (10) Clark, G; Nucleic Acids Res 1997, V25, P1510 HCAPLUS
- (11) Czarny, A; J Am Chem Soc 1995, V117, P4716 HCAPLUS
- (12) Drew, H; Cell 1984, V37, P491 HCAPLUS
- (13) Drobyshev, A; Nucleic Acids Res 1999, V27, P4100 HCAPLUS
- (14) Ebrahimi, S; J Chem Soc, Chem Commun 1992, P1398 HCAPLUS
- (15) Embrey, K; Eur J Biochem 1993, V211, P437 HCAPLUS
- (16) Embrey, K; J Chem Soc, Chem Commun 1991, P1770 HCAPLUS
- (17) Fede, A; Biochemistry 1991, V30, P11377 HCAPLUS
- (18) Fede, A; Structure 1993, V1, P177 HCAPLUS
- (19) Gavathiotis, E; Nucleic Acids Res 2000, V28, P728 HCAPLUS
- (20) Gerber, P; J Comput Aided Mol Des 1995, V9, P251 HCAPLUS
- (21) Grimmet, M; Compressive Heterocyclic Chemistry 1984, V5, P457
- (22) Haq, I; J Mol Biol 1997, V271, P244 HCAPLUS
- (23) Hauglang, R; Handbook of Fluorescent Probes and Research Chemicals 6th ed
1996
- (24) Jorgenson, K; J Biomol Struct Dyn 1988, V5, P1005 HCAPLUS
- (25) Kopka, M; Nucleic Acids and Molecular Biology 1987, V1, P1
- (26) Kumar, S; J Biomol Struct Dyn 1992, V8, P331
- (27) Loontjens, F; Biochemistry 1990, V29, P9029 HCAPLUS
- (28) Loontjens, F; Biochemistry 1991, V30, P182 HCAPLUS
- (29) Low, C; Nucleic Acids Res 1984, V12, P4865 HCAPLUS
- (30) Maquestiau, A; Synthetic Commun 1991, V21, P2181
- (31) Nahm, S; Tetrahedron Lett 1981, V22, P3815 HCAPLUS
- (32) Parkinson, J; Biochemistry 1990, V29, P10181 HCAPLUS
- (33) Parkinson, J; Biochemistry 1994, V33, P8442 HCAPLUS
- (34) Pilch, D; Drug Des Discov 1996, V13, P115 HCAPLUS
- (35) Pjura, P; J Mol Biol 1987, V197, P257 HCAPLUS
- (36) Portugal, J; Biochim Biophys Acta 1987, V949, P158


- (37) Quintana, J; Biochemistry 1991, V30, P10294 HCAPLUS
 (38) Rao, K; Chem Res Toxicol 1991, V4, P661 HCAPLUS
 (39) Satz, A; J Am Chem Soc 2001, V123, P2469 HCAPLUS
 (40) Searle, M; Nucleic Acids Res 1990, V18, P3753 HCAPLUS
 (41) Sigman, D; Annu Rev Biochem 1990, V59, P207 HCAPLUS
 (42) Singh, M; Chem Res Toxicol 1992, V5, P597 HCAPLUS
 (43) Spink, N; Nucleic Acids Res 1994, V22, P1607 HCAPLUS
 (44) Sriram, M; EMBO J 1992, V11, P225 HCAPLUS
 (45) Starshikov, N; Khim Chem Heterocycl Comp 1980, P81
 (46) Stokke, T; J Histochem Cytochem 1985, V33, P333 HCAPLUS
 (47) Suck, D; Nature 1988, V332, P465
 (48) Thurston, D; Cancer 1999, V80, P65 HCAPLUS
 (49) Van Dyke, M; Biochemistry 1983, V22, P2373 HCAPLUS
 (50) Vega, M; Eur J Biochem 1994, V222, P721 HCAPLUS
 (51) Vogt, H; Rev Sci Instrum 1995, V66, P4385 HCAPLUS
 (52) Von Loewe, H; Arzneim Forsch 1974, V24, P1927
 (53) Weston, S; J Mol Biol 1992, V226, P1237 HCAPLUS
 (54) Wood, A; Nucleic Acids Res 1995, V23, P3678 HCAPLUS

IT 182496-21-5P 391903-25-6P

RL: SPN (Synthetic preparation); PREP (Preparation)
 (design, synthesis and DNA sequence recognition using
 tris-benzimidazole derivs.)


RN 182496-21-5 HCAPLUS

CN 3-Pyrrolidinamine, 1-[2'-(4-methoxyphenyl)[2,5':2',5''-ter-1H-benzimidazol]-5-yl]- (9CI) (CA INDEX NAME)



RN 391903-25-6 HCAPLUS

CN 2,5':2',5''-Ter-1H-benzimidazole, 2'-(4-methoxyphenyl)-5-(4-methyl-1-piperazinyl)- (9CI) (CA INDEX NAME)



L28 ANSWER 5 OF 17 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 2001:656650 HCAPLUS

DN 135:368118

ED Entered STN: 07 Sep 2001

TI Molecular modelling of ligand-DNA minor groove binding: role of
 ligand-water interactions

AU Mikheikin, A. L.; Zhuze, A. L.; Zasedatelev, A. S.

CS Engelhardt Institute of Molecular Biology, Russian Academy of Sciences,
 Moscow, 119991, Russia

SO Journal of Biomolecular Structure & Dynamics (2001), 19(1), 175-178

CODEN: JBSDD6; ISSN: 0739-1102

PB Adenine Press

DT Journal


LA English

CC 6-2 (General Biochemistry)

AB A procedure was developed for quant. estimation of the ligand affinity for the
 DNA minor groove with allowance for ligand hydration, whereby the binding
 energy was calculated as the difference in the energies of ligand-DNA and
 ligand-water interactions. Adequacy of the procedure was demonstrated
 with the structural motifs (pyrrolecarboxamide, benzimidazole,
 furancarboxamide, and phthalimide) of well-known ligands for the case of a
 d(GCA10CG).cntdot.d(CGT10GC) duplex. On the strength of the results

obtained, an indole-based motif was proposed as the basis for a highly
affined minor groove binder.

ST DNA minor groove interaction ligand
IT Hydration, chemical
(hydration and hydrophobic interactions in ligand-DNA complexes also
play role in binding of ligand to DNA minor groove)
IT Molecular association
(mol. modeling of ligand-DNA minor groove binding and role of
ligand-water interactions)
IT DNA
RL: BPR (Biological process); BSU (Biological study, unclassified); PRP
(Properties); BIOL (Biological study); PROC (Process)
(mol. modeling of ligand-DNA minor groove binding and role of
ligand-water interactions)
IT 7732-18-5, Water, biological studies
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL
(Biological study); PROC (Process)
(mol. modeling of ligand-DNA minor groove binding and role of
ligand-water interactions)
IT 636-47-5, Distamycin A 23491-45-4, Hoechst 33258 47165-04-8, DAPI
168100-52-5 373596-17-9 373596-18-0 373596-19-1
373596-20-4 373596-21-5 373596-22-6
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(mol. modeling of ligand-DNA minor groove binding and role of
ligand-water interactions)
RE.CNT 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD
RE
(1) Bostock-Smith, C; Nucleic Acids Res 1999, V27, P1619 HCAPLUS
(2) Haq, I; J Mol Biol 1997, V271, P244 HCAPLUS
(3) Kopka, M; J Mol Biol 1985, V183, P553 HCAPLUS
(4) Kopka, M; Proc Natl Acad Sci USA 1985, V82, P1376 HCAPLUS
(5) Kubista, M; Biochemistry 1987, V26, P4545 HCAPLUS
(6) Mikhailov, M; Mol Biol (Moscow) 1981, V15, P690 HCAPLUS
(7) Mikheikin, A; Mol Biol (Moscow) 1997, V31, P854 HCAPLUS
(8) Salmanova, D; Mol Biol (Moscow) 1995, V29, P491
(9) Yu, N; Biopolymers 1995, V35, P377
(10) Zasedatelev, A; Dokl Akad Nauk SSSR 1976, V231, P1006 HCAPLUS
IT 373596-19-1
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(mol. modeling of ligand-DNA minor groove binding and role of
ligand-water interactions)
RN 373596-19-1 HCAPLUS
CN [2,5':2',5''-Ter-1H-benzimidazole]-2'',5-diamine (9CI) (CA INDEX NAME)



L28 ANSWER 6 OF 17 HCAPLUS COPYRIGHT 2005 ACS on STN
AN 2000:748488 HCAPLUS
DN 134:291615
ED Entered STN: 24 Oct 2000
TI Molecular recognition of DNA by Hoechst benzimidazoles: exploring beyond
the pyrrole-imidazole-hydroxypyrrole polyamide-pairing code
AU Minehan, Thomas G.; Gottwald, Konstanze; Dervan, Peter B.
CS Division of Chemistry and Chemical Engineering, California Institute of
Technology, Pasadena, CA, 91125, USA
SO Helvetica Chimica Acta (2000), 83(9), 2197-2213
CODEN: HCACAV; ISSN: 0018-019X
PB Verlag Helvetica Chimica Acta
DT Journal
LA English
CC 6-2 (General Biochemistry)
Section cross-reference(s): 28
OS CASREACT 134:291615
AB A series of three-ring analogs of the minor-groove-binding mol. Hoechst
33258 (1), consisting of benzimidazole (B), imidazopyridine (P), and
hydroxybenzimidazole (H) monomers, have been synthesized in order to

investigate both their sequence specificity and binding modes. MPE.cntdot.FeII Footprinting has revealed the preference of both PBB and BBB ligands for 5'-WGWWW-3' and 5'-WCWWW-3' tracts, as well as A.cntdot.T-rich sequences. Affinity-cleavage titrns. show no evidence for a 2:1 binding mode of these Hoechst analogs. Importantly, all derivs. are oriented in one direction at each of their binding sites. The implications of these results for the design of minor-groove-binding small mols. is discussed.

ST DNA recognition Hoechst 33258 benzimidazole analog prepn
IT DNA
RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)
(double-stranded; preparation and mol. recognition of DNA minor groove by Hoechst 33258 benzimidazole analogs)
IT Molecular orientation
(in minor-groove; preparation and mol. recognition of DNA minor groove by Hoechst 33258 benzimidazole analogs)
IT Molecular recognition
(preparation and mol. recognition of DNA minor groove by Hoechst 33258 benzimidazole analogs)
IT 23491-45-4, Hoechst 33258 334685-20-0 334685-35-7
RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)
(preparation and mol. recognition of DNA minor groove by Hoechst 33258 benzimidazole analogs)
IT 334685-29-9P 334685-30-2P 334685-33-5P 334685-34-6P
RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); PROC (Process)
(preparation and mol. recognition of DNA minor groove by Hoechst 33258 benzimidazole analogs)
IT 105-36-2, Ethyl bromoacetate 105-83-9, Bis(3-aminopropyl)methylamine 109-55-7, 3-(Dimethylamino)propylamine 110-85-0, Piperazine, reactions 1635-61-6, 5-Chloro-2-nitroaniline 6291-84-5, (3-Aminopropyl)methylamine 23911-25-3, EDTA dianhydride 126463-85-2 126824-22-4 142764-79-2 183296-71-1
RL: RCT (Reactant); RACT (Reactant or reagent)
(preparation and mol. recognition of DNA minor groove by Hoechst 33258 benzimidazole analogs)
IT 96103-52-5P 126436-26-8P 126463-87-4P 167959-17-3P 188247-41-8P 188247-43-0P 334685-22-2P 334685-25-5P 334685-27-7P 334685-31-3P 334685-32-4P 416850-41-4P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and mol. recognition of DNA minor groove by Hoechst 33258 benzimidazole analogs)
IT 334685-21-1P 334685-28-8P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation and mol. recognition of DNA minor groove by Hoechst 33258 benzimidazole analogs)
RE.CNT 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD
RE
(1) Coll, M; Proc Natl Acad Sci 1987, V84, P8385 HCAPLUS
(2) Czarny, A; J Heterocycl Chem 1996, V33, P1393 HCAPLUS
(3) Dervan, P; Curr Opin Chem Biol 1999, V3, P688 HCAPLUS
(4) Dervan, P; Science 1986, V232, P464 HCAPLUS
(5) Frau, S; Bull Soc Chim Fr 1996, V133, P1053 HCAPLUS
(6) Harshman, K; Nucl Acids Res 1985, V13, P4825 HCAPLUS
(7) Herman, D; J Am Chem Soc 1998, V120, P1382 HCAPLUS
(8) Klevit, R; Biochemistry 1986, V25, P3296 HCAPLUS
(9) Kopka, M; J Mol Biol 1985, V183, P553 HCAPLUS
(10) Kopka, M; Proc Natl Acad Sci 1985, V82, P1376 HCAPLUS
(11) Kubista, M; Biochemistry 1987, V26, P4545 HCAPLUS
(12) Kumar, S; J Biomol Struct Dyn 1990, V8, P331 HCAPLUS
(13) Lee, M; Heterocycles 1991, V32, P2093 HCAPLUS
(14) Lown, J; Advances in DNA Sequence Specific Agents 1998, V3, P67 HCAPLUS
(15) Martin, R; Nature 1983, V302, P452 HCAPLUS
(16) Mrksich, M; Proc Natl Acad Sci 1992, V89, P7568
(17) Pelton, J; Biochemistry 1988, V27, P8088 HCAPLUS
(18) Pelton, J; J Am Chem Soc 1990, V112, P1393 HCAPLUS
(19) Pelton, J; Proc Natl Acad Sci 1989, V86, P5723 HCAPLUS
(20) Pjura, P; J Mol Biol 1987, V197, P257 HCAPLUS
(21) Schrodinger Inc; Jaguar software package
(22) Singh, M; Chem Res Toxicol 1992, V5, P597 HCAPLUS
(23) Tanaka, A; Chem Pharm Bull 1994, V42, P560 HCAPLUS
(24) Teng, M; Nucl Acids Res 1988, V16, P2671 HCAPLUS


- (25) Trauger, J; J Am Chem Soc 1996, V118, P6160 HCAPLUS
 (26) Van Dyke, M; Nucl Acids Res 1983, V11, P5555 HCAPLUS
 (27) Wade, W; J Am Chem Soc 1992, V114, P8783 HCAPLUS
 (28) Wang, L; Proc Natl Acad Sci 2000, V97, P12 HCAPLUS
 (29) Wemmer, D; Curr Opin Struct Biol 1997, V7, P355 HCAPLUS
 (30) Zimmer, C; Prog Biophys Mol Biol 1986, V47, P31 HCAPLUS

IT 334685-29-9P 334685-33-5P

RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); PROC (Process)
 (preparation and mol. recognition of DNA minor groove by Hoechst 33258 benzimidazole analogs)

RN 334685-29-9 HCAPLUS


CN [2,5':2',5''-Ter-1H-benzimidazole]-5-carboxamide, N-[3-(dimethylamino)propyl]-2'',3''-dihydro-2''-oxo- (9CI) (CA INDEX NAME)




RN 334685-33-5 HCAPLUS

CN 2,6,10,13,16-Pentaazaoctadecan-18-oic acid, 13,16-bis(carboxymethyl)-1-(2'',3''-dihydro-2''-oxo[2,5':2',5''-ter-1H-benzimidazol]-5-yl)-6-methyl-1,11-dioxo- (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B




IT 167959-17-3P 334685-27-7P 334685-31-3P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (preparation and mol. recognition of DNA minor groove by Hoechst 33258 benzimidazole analogs)

RN 167959-17-3 HCAPLUS

CN [2,5':2',5''-Ter-1H-benzimidazole]-5-carbonitrile, 2''-(4-methoxyphenyl)- (9CI) (CA INDEX NAME)




RN 334685-27-7 HCAPLUS
 CN [2,5':2',5''-Ter-1H-benzimidazole]-5-carbonitrile, 7''-chloro-4''-methoxy-2''-methyl- (9CI) (CA INDEX NAME)



RN 334685-31-3 HCAPLUS
 CN [2,5':2',5''-Ter-1H-benzimidazole]-5-carboxamide, N-[3-[(3-aminopropyl)methylamino]propyl]-2'',3''-dihydro-2''-oxo- (9CI) (CA INDEX NAME)


PAGE 1-A




PAGE 1-B

=O


IT 334685-21-1P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation and mol. recognition of DNA minor groove by Hoechst 33258
 benzimidazole analogs)
 RN 334685-21-1 HCAPLUS
 CN [2,5':2',5''-Ter-1H-benzimidazole]-5-carboxamide, 7''-chloro-N-[3-(dimethylamino)propyl]-4''-hydroxy-2''-methyl- (9CI) (CA INDEX NAME)



L28 ANSWER 7 OF 17 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 2000:124956 HCAPLUS
 DN 132:274285
 ED Entered STN: 24 Feb 2000
 TI The interaction of benzimidazole compounds with DNA: intercalation and groove binding modes
 AU Kubota, Yukio; Iwamoto, Takayuki; Seki, Toshimasa
 CS Department of Chemistry, Faculty of Science, Yamaguchi University, Yamaguchi, 753-8512, Japan
 SO Nucleic Acids Symposium Series (1999), 42 (Twenty-sixth Symposium on Nucleic Acids Chemistry, 1999), 53-54
 CODEN: NACSD8; ISSN: 0261-3166
 PB Oxford University Press
 DT Journal
 LA English
 CC 1-12 (Pharmacology)
 AB Benzimidazole compds. have been synthesized to study their DNA-binding properties. Results obtained with spectroscopy and viscosity measurements indicate that the binding mode varies from intercalation to groove-binding, depending on the number of benzimidazole rings (conformation and size of compds.).
 ST benzimidazole intercalation DNA intercalation
 IT Conformation
 Molecular association
 (interaction of benzimidazole compds. with DNA in relation to intercalation and groove binding modes and conformation)
 IT DNA
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
 (interaction of benzimidazole compds. with DNA in relation to intercalation and groove binding modes and conformation)
 IT Intercalation
 (nucleic acid; interaction of benzimidazole compds. with DNA in relation to intercalation and groove binding modes and conformation)
 IT 154713-23-2 263707-95-5 263707-96-6 263707-97-7
 RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)
 (interaction of benzimidazole compds. with DNA in relation to intercalation and groove binding modes and conformation)
 RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD
 RE
 (1) Bathini, Y; Chem Res Toxicol 1990, V3, P268 HCAPLUS
 (2) Kubota, K; Biochim Biophys Acta 1977, V478, P23
 (3) Kubota, Y; Chem Lett 1991, P745 HCAPLUS
 (4) McGhee, J; J Mol Biol 1974, V86, P469 HCAPLUS
 (5) Nakano, S; Nucleic Acids Sym Ser 1994, V31, P75 HCAPLUS
 (6) Zimmer, C; Prog Biophys Mol Biol 1986, V47, P31 HCAPLUS
 IT 263707-96-6 263707-97-7
 RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)
 (interaction of benzimidazole compds. with DNA in relation to intercalation and groove binding modes and conformation)
 RN 263707-96-6 HCAPLUS
 CN 2,5':2',5''-Ter-1H-benzimidazole, 5-(4-methyl-1-piperazinyl)- (9CI) (CA INDEX NAME)




RN 263707-97-7 HCAPLUS
 CN Phenol, 4-[5-(4-methyl-1-piperazinyl)[2,5':2',5''-ter-1H-benzimidazol]-2''-yl]- (9CI) (CA INDEX NAME)



L28 ANSWER 8 OF 17 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 1999:446949 HCAPLUS
 DN 132:60607
 ED Entered STN: 21 Jul 1999
 TI DNA minor groove recognition of a non-self-complementary AT-rich sequence by a tris-benzimidazole ligand
 AU Aymami, Juan; Nunn, Christine M.; Neidle, Stephen
 CS CRC Biomolecular Structure Unit, Institute of Cancer Research, Surrey, SM2 5NG, UK
 SO Nucleic Acids Research (1999), 27(13), 2691-2698
 CODEN: NARHAD; ISSN: 0305-1048
 PB Oxford University Press
 DT Journal
 LA English
 CC 6-2 (General Biochemistry)
 Section cross-reference(s): 1, 75
 AB The crystal structure of the non-self-complementary dodecamer DNA duplex formed by d(CG[5BrC]ATATTGCG) and d(CGCAAATATGCG) has been solved to 2.3 .ANG. resolution, together with that of its complex with the tris-benzimidazole minor groove binding ligand TRIBIZ. The inclusion of a bromine atom on one strand in each structure enabled the possibility of disorder to be discounted. The native structure has an exceptional narrow minor groove, of 2.5-2.6 .ANG. in the central part of the A/T region, which is increased in width by .apprx.0.8 .ANG. on drug binding. The ligand mol. binds in the central part of the sequence. The benzimidazole subunits of the ligand participate in six bifurcated hydrogen bonds with A:T base pair edges, three to each DNA strand. The presence of a pair of C-H...O hydrogen bonds has been deduced from the close proximity of the pyrrolidine group of the ligand to the Tpa step in the sequence.
 ST DNA TRIBIZ structure recognition
 IT Crystal structure
 Molecular recognition
 (DNA minor groove recognition of non-self-complementary AT-rich sequence by tris-benzimidazole ligand)
 IT Conformation
 (DNA; DNA minor groove recognition of non-self-complementary AT-rich sequence by tris-benzimidazole ligand)
 IT Molecular structure
 (cDNA minor groove recognition of non-self-complementary AT-rich sequence by tris-benzimidazole ligand)
 IT 182496-21-5, TRIBIZ 253331-47-4
 RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)
 (DNA minor groove recognition of non-self-complementary AT-rich sequence by tris-benzimidazole ligand)
 RE.CNT 48 THERE ARE 48 CITED REFERENCES AVAILABLE FOR THIS RECORD
 RE
 (1) Abu-Daya, A; Nucleic Acids Res 1995, V23, P3385 HCAPLUS
 (2) Bostock-Smith, C; Nucleic Acids Res 1999, V27, P1619 HCAPLUS
 (3) Brown, D; J Mol Biol 1992, V226, P481 HCAPLUS
 (4) Brunger, A; Science 1987, V235, P458
 (5) Chen, X; J Mol Biol 1997, V267, P1157 HCAPLUS
 (6) Clark, G; Biochemistry 1996, V35, P13745 HCAPLUS
 (7) Coll, M; Proc Natl Acad Sci 1987, V84, P8385 HCAPLUS
 (8) DiGabriele, A; J Mol Biol 1993, V231, P1024 HCAPLUS
 (9) DiGabriele, A; Proc Natl Acad Sci 1989, V86, P1816 HCAPLUS
 (10) Dickerson, R; Proc Natl Acad Sci 1994, V91, P3579 HCAPLUS
 (11) Dickinson, L; Proc Natl Acad Sci 1998, V95, P12890 HCAPLUS
 (12) Edwards, K; J Mol Biol 1992, V226, P1161 HCAPLUS
 (13) Embrey, K; Eur J Biochem 1993, V211, P437 HCAPLUS
 (14) Goodsell, D; Biochemistry 1995, V34, P16654 HCAPLUS
 (15) Goodsell, D; Biochemistry 1995, V34, P4983 HCAPLUS
 (16) Gottesfield, J; Nature 1997, V387, P202

- (17) Guerri, A; Nucleic Acids Res 1998, V26, P2873 HCAPLUS
 (18) Kielkopf, C; Nature Struct Biol 1998, V5, P104 HCAPLUS
 (19) Kielkopf, C; Science 1998, V282, P111 HCAPLUS
 (20) Kim, J; Bioorg Med Chem 1998, V6, P163 HCAPLUS
 (21) Kopka, M; Structure 1997, V5, P1033 HCAPLUS
 (22) Lavery, R; J Biomol Struct Dyn 1989, V6, P655 HCAPLUS
 (23) Neidle, S; Biopolymers 1997, V44, P105 HCAPLUS
 (24) Nelson, H; Nature 1987, V330, P221 HCAPLUS
 (25) Nunn, C; Biochemistry 1997, V36, P4792 HCAPLUS
 (26) Otwinowski, Z; Data Collection and Processing 1993
 (27) Parkinson, G; Acta Crystallogr 1996, V52, P57 HCAPLUS
 (28) Pilch, D; Proc Natl Acad Sci 1997, V94, P13565 HCAPLUS
 (29) Rentzeperis, D; Biochemistry 1995, V34, P2937 HCAPLUS
 (30) Rydzewski, J; Nucleic Acids Res 1996, V24, P1287 HCAPLUS
 (31) Sheldrick, G; SHELX a Crystallographic Refinement Programme 1997
 (32) Shui, X; Biochemistry 1998, V37, P8341 HCAPLUS
 (33) Singh, S; J Am Chem Soc 1999, V121, P3267 HCAPLUS
 (34) Spink, N; Nucleic Acids Res 1994, V22, P1607 HCAPLUS
 (35) Squire, C; Nucleic Acids Res 1997, V25, P4072 HCAPLUS
 (36) Trauger, J; Nature 1996, V382, P559 HCAPLUS
 (37) Trent, J; J Med Chem 1996, V39, P4554 HCAPLUS
 (38) Vega, M; Eur J Biochem 1994, V222, P721 HCAPLUS
 (39) Vega, M; Eur J Biochem 1996, V239, P376 HCAPLUS
 (40) Walker, W; Proc Natl Acad Sci 1997, V94, P5634 HCAPLUS
 (41) Walker, W; Proc Natl Acad Sci 1998, V95, P4315 HCAPLUS
 (42) White, S; Chem Biol 1997, V4, P569 HCAPLUS
 (43) White, S; Nature 1998, V391, P468 HCAPLUS
 (44) Wood, A; Nucleic Acids Res 1995, V23, P3678 HCAPLUS
 (45) Xu, Z; Biochemistry 1998, V37, P3558 HCAPLUS
 (46) Yoon, C; Proc Natl Acad Sci 1988, V85, P6332 HCAPLUS
 (47) Yuan, H; Biochemistry 1992, V31, P8009 HCAPLUS
 (48) Zimmer, C; Prog Biophys Mol Biol 1986, V47, P31 HCAPLUS
 IT 182496-21-5, TRIBIZ
 RL: BPR (Biological process); BSU (Biological study, unclassified); PRP
 (Properties); BIOL (Biological study); PROC (Process)
 (DNA minor groove recognition of non-self-complementary AT-rich
 sequence by tris-benzimidazole ligand)
 RN 182496-21-5 HCAPLUS
 CN 3-Pyrrolidinamine, 1-[2'-(4-methoxyphenyl)[2,5':2',5''-ter-1H-
 benzimidazol]-5-yl]- (9CI) (CA INDEX NAME)



- L28 ANSWER 9 OF 17 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 1998:523091 HCAPLUS
 DN 129:269959
 ED Entered STN: 21 Aug 1998
 TI Ligands for DNA and RNA
 AU Douglas, Kenneth T.
 CS School of Pharmacy and Pharmaceutical Sciences, University of Manchester,
 Manchester, M13 9PL, UK
 SO Structure, Motion, Interaction and Expression of Biological
 Macromolecules, Proceedings of the Conversation in the Discipline
 Biomolecular Stereodynamics, 10th, Albany, June 17-21, 1997 (1998),
 Meeting Date 1997, Volume 1, 279-293. Editor(s): Sarma, Ramaswamy H.;
 Sarma, Mukti H. Publisher: Adenine Press, Schenectady, N. Y.
 CODEN: 66NGAV
 DT Conference
 LA English
 CC 1-3 (Pharmacology)
 AB In contrast to rational drug and ligand design based on mol. graphics for
 the field of proteins, the situation for nucleic acid ligand design is
 less advanced, but is now proceeding rapidly. For DNA the aspect most
 developed in this sense is the minor groove, but for RNA there is still
 rather little in the way of ligand design reported. Insight into the DNA
 minor groove has reached the stage at which it is now possible to test the

possibility of rational ligand design in a number of objective ways. The first stage of this is to rationalize observations already made, but more rigorous is to test the ability to predict structural and chemical properties. In this contribution the ability to predict binding interactions for ligands in the minor groove of B-DNA of a series of analogs of Hoechst 33258 will be analyzed. These compds. were designed using mol. graphic/dynamics based on high-field NMR structural determination of Hoechst-duplex DNA complexes using a synthetic oligonucleotide sequence. Enthalpy and entropy contributors to net binding strength will be considered. The test of prediction powers is not merely to be able to achieve better net ligand binding strength, but also to propose specific bonding interactions. These predictions have been probed structurally using NMR anal. at high resolution of ligand-DNA complexes, again designed by mol. modeling. As well as using structural probes such as NMR spectroscopy, it is possible to test predictive ability by introducing novel chemical reactivity. In this context we shall describe a novel DNA strand-cleaving method, designed using mol. graphics of the above structures to locate a transition metal ion binding site very specifically and close to the phosphodiester backbone, allowing the generation of reactive free radicals to effect cleavage. Relative to DNA, RNA-binding ligands are less widely studied at present and, in the final part of the contribution, the binding properties of some new ligands for tRNA, binding with 1:1 stoichiometry and low micromolar dissociation consts. will be described. Their binding has been studied by UV-visible spectrophotometry, fluorescent titration and NMR spectroscopy.

ST DNA RNA ligand structure modeling
 IT Simulation and Modeling, biological
 Structure-activity relationship
 (mol. modeling of ligands for DNA and RNA)
 IT Ligands
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)
 (mol. modeling of ligands for DNA and RNA)
 IT DNA
 RNA
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
 (mol. modeling of ligands for DNA and RNA)
 IT 23491-45-4, Hoechst 33258 23491-53-4 39389-47-4, Distamycin
 90991-94-9 132869-83-1 171782-32-4 171782-33-5 213974-59-5
 213974-61-9
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)
 (mol. modeling of ligands for DNA and RNA)

RE.CNT 75 THERE ARE 75 CITED REFERENCES AVAILABLE FOR THIS RECORD
 RE

- (1) Baindur, S; Biochim Biophys Acta 1987, V923, P66 HCAPLUS
- (2) Baindur, S; J Chem Soc Chem Commun 1985, P1398 HCAPLUS
- (3) Birdsall, W; Biochim Biophys Acta 1989, V1007, P176 HCAPLUS
- (4) Borisova, O; Mol Biol (USA) 1974, V7, P411 MEDLINE
- (5) Brown, R; Nature (London) 1983, V303, P543 HCAPLUS
- (6) Bunni, M; Biochem J 1984, V217, P383 HCAPLUS
- (7) Bunni, M; Biochim Biophys Acta 1987, V923, P421 HCAPLUS
- (8) Carter, B; Proc Nat Acad Sci (U S A) 1990, V87, P9373 HCAPLUS
- (9) Celander, D; Biochemistry 1996, V35, P12061 HCAPLUS
- (10) Chow, C; J Am Chem Soc 1990, V112, P2839 HCAPLUS
- (11) Ciesiolka, J; Eur J Biochem 1989, V186, P71 HCAPLUS
- (12) Clark, G; Nucl Acids Res 1996, V24, P4882 HCAPLUS
- (13) Dange, V; Science 1990, V248, P585 HCAPLUS
- (14) de Carrondo, M; Biochemistry 1989, V28, P7849
- (15) Douglas, K; Progress in Bio-organic Chemistry 1976, P194
- (16) Ebrahimi, S; Anti-Cancer Drug Design 1995, V10, P463 HCAPLUS
- (17) Ebrahimi, S; J Chem Soc Chem Commun 1992, P1398 HCAPLUS
- (18) Embrey, K; Eur J Biochem 1993, V211, P437 HCAPLUS
- (19) Embrey, K; Eur J Biochem 1994, V221, P793 HCAPLUS
- (20) Embrey, K; J Chem Soc Chem Commun 1991, P1770 HCAPLUS
- (21) Famulok, M; Biochemistry 1996, V35, P4265 HCAPLUS
- (22) Fede, A; Biochemistry 1991, V30, P11377 HCAPLUS
- (23) Fede, A; Structure 1993, V1, P177 HCAPLUS
- (24) Pontecilla-Camps, J; J Am Chem Soc 1979, V101, P6114 HCAPLUS
- (25) Poster, N; Biochim Biophys Acta 1988, V950, P118 HCAPLUS
- (26) Frydman, L; Proc Natl Acad Sci USA 1992, V89, P9186 HCAPLUS
- (27) Goodsell, D; J Med Chem 1986, V29, P727 HCAPLUS
- (28) Harshman, K; Nucl Acids Res 1985, V13, P4825 HCAPLUS
- (29) Heerschap, A; Nucl Acids Res 1986, V14, P983 HCAPLUS
- (30) Heerschap, A; Nucl Acids Res 1986, V14, P983 HCAPLUS


- (31) Heitner, H; J Am Chem Soc 1972, V94, P8936 HCAPLUS
 (32) Hutlin, T; Chem -Biol Interact 1974, V8, P315
 (33) Jin, R; Proc Natl Acad Sci USA 1988, V85, P8939 HCAPLUS
 (34) Kazagov, S; Proc Nat Acad Sci (U S A) 1992, V89, P7939
 (35) Kean, J; Biochemistry 1985, V24, P5062 HCAPLUS
 (36) Kopka, M; Proc Natl Acad Sci USA 1985, V82, P1376 HCAPLUS
 (37) Leupin, W; Actual Chim Ther 1994, V21, P153 HCAPLUS
 (38) Loontjens, F; Biochemistry 1990, V29, P9029 HCAPLUS
 (39) Loontjens, F; Biochemistry 1991, V30, P182 HCAPLUS
 (40) Matthews, D; J Cell Biochem 1978, V253, P6946 HCAPLUS
 (41) Mitsuya, H; Science 1990, V249, P1533 HCAPLUS
 (42) Mrksich, M; J Am Chem Soc 1994, V116, P3663 HCAPLUS
 (43) Murakawa, G; Nucl Acids Res 1989, V17, P5361 HCAPLUS
 (44) Nikolaev, V; J Biomol Struct Dyn 1996, V14, P31 HCAPLUS
 (45) Page, M; Proc Natl Acad Sci USA 1971, V68, P1678 HCAPLUS
 (46) Pal, B; Arch Biochem Biophys 1972, V150, P86 HCAPLUS
 (47) Parkinson, J; Biochemistry 1990, V29, P10181 HCAPLUS
 (48) Parkinson, J; Biochemistry 1994, V33, P8442 HCAPLUS
 (49) Parkinson, J; Biochemistry 1995, V34, P16240 HCAPLUS
 (50) Parkinson, J; J Chem Soc Chem Commun 1989, P1023 HCAPLUS
 (51) Parkinson, J; Mag Res Chem 1992, V30, P1064 HCAPLUS
 (52) Peterson, R; J Mol Biol 1996, V264, P863 HCAPLUS
 (53) Pilch, D; Biochemistry 1995, V34, P16107 HCAPLUS
 (54) Pilch, D; Biochemistry 1995, V34, P9962 HCAPLUS
 (55) Pilch, D; Proc Natl Acad Sci USA 1996, V93, P8306 HCAPLUS
 (56) Pjura, P; J Mol Biol 1987, V197, P257 HCAPLUS
 (57) Quintana, J; Biochemistry 1991, V30, P10294 HCAPLUS
 (58) Rao, K; Anti-Cancer Drug Design 1990, V5, P3 HCAPLUS
 (59) Ratmeyer, L; Biochemistry 1996, V35, P13689 HCAPLUS
 (60) Rubin, J; J Biomol Struct Dynamics 1984, V2, P165 HCAPLUS
 (61) Sadat-Ebrahimi, S; J Chem Soc Chem Commun 1997, P385
 (62) Searle, M; Nucl Acids Res 1990, V18, P3753 HCAPLUS
 (63) Surovaya, A; J Biomol Struct Dyn 1997, V14, P1
 (64) Surovaya, A; Mol Biol (USA) 1974, V7, P403 MEDLINE
 (65) Tanious, F; Biochemistry 1992, V31, P3103 HCAPLUS
 (66) Teng, M; Nucl Acids Res 1988, V16, P2671 HCAPLUS
 (67) Totusuka, R; Biochem Biophys Res Commun 1995, V208, P168 HCAPLUS
 (68) Urbanke, C; Eur J Biochem 1973, V33, P511 HCAPLUS
 (69) Walba, H; J Org Chem 1961, V26, P2789 HCAPLUS
 (70) Wallis, M; Chemistry and Biology 1995, V2, P543 HCAPLUS
 (71) Wang, Y; Chemistry and Biology 1995, V2, P281 HCAPLUS
 (72) Westhof, E; Biochemistry 1986, V25, P4868 HCAPLUS
 (73) Wilson, W; Biochemistry 1993, V32, P4098 HCAPLUS
 (74) Zakrzewska, K; J Biomol Struct Dyn 1984, V2, P737
 (75) Zimmer, C; J Mol Biol 1971, V58, P329 HCAPLUS

IT 23491-53-4

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)
 (mol. modeling of ligands for DNA and RNA)

RN 23491-53-4 HCAPLUS

CN 2,5'-Bi-1H-benzimidazole, 2'-(1,3-benzodioxol-5-yl)-5-(4-methyl-1-piperazinyl)- (9CI) (CA INDEX NAME)



L28 ANSWER 10 OF 17 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1998:512965 HCAPLUS

DN 129:239452

ED Entered STN: 19 Aug 1998

TI Molecular modeling and footprinting studies of DNA minor groove binders: bisquaternary ammonium heterocyclic compounds

AU Slickers, P.; Hillebrand, M.; Kittler, L.; Lober, G.; Suhnel, J.

CS Inst. Mol. Biotechnol., Jena, D-07708, Germany

SO Anti-Cancer Drug Design (1998), 13(5), 463-488

CODEN: ACDDEA; ISSN: 0266-9536

PB Oxford University Press

DT Journal

LA English

CC 1-3 (Pharmacology)
 Section cross-reference(s): 22

AB The authors report new quant. footprinting data which reveal differences in binding consts. of bisquaternary ammonium heterocyclic compds. (BQA) with AT-rich DNA sites depending on the ligand structure and on the size and sequence of the DNA binding site. To understand the dependence of binding affinity on the ligand structure the authors have performed quantum-chemical AM1 calcns. on the BQA compds. and on subunits to explore the conformational space and to calculate the electronic and structural features of individual ligand conformations. Due to the properties of the rotatable backbone bonds, there is a large number of possible conformations with almost equal energy. The authors present a new method for the calcn. of the radius of curvature of mol. structures. Assuming that strong binders should have a shape complementary to the DNA minor groove, this measure is used to select the optimum conformations for DNA-drug binding. The approach yields the correct ligand conformation for SN6999, for which an x-ray DNA-drug structure is known. The curvature of the optimum conformations of all ligands is compared with the exptl. binding consts. A correlation is found between curvature and binding constant provided other structural factors do not vary. Therefore, the authors conclude that within structurally similar BQA compds., the extent of curvature is the relevant quantity which modulates the binding affinity.

ST DNA minor groove binder heterocyclic compd; bisquaternary ammonium heterocyclic compd DNA binding; mol modeling DNA binding heterocyclic compd; QSAR DNA binding heterocyclic compd

IT Bond angle
 (dihedral; mol. modeling and footprinting studies of DNA minor groove binders using bisquaternary ammonium heterocyclic compds.)

IT AM1 MO (molecular orbital)
 Bond length
 Conformation
 Electrostatic potential
 Molecular association
 Molecular modeling
 QSAR (structure-activity relationship)
 (mol. modeling and footprinting studies of DNA minor groove binders using bisquaternary ammonium heterocyclic compds.)

IT DNA
 RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)
 (mol. modeling and footprinting studies of DNA minor groove binders using bisquaternary ammonium heterocyclic compds.)

IT Bond angle
 (torsional; mol. modeling and footprinting studies of DNA minor groove binders using bisquaternary ammonium heterocyclic compds.)

IT 14120-88-8, SN4094 18355-40-3, SN6570 23491-45-4, Hoechst 33258
 23617-49-4, SN6324 23647-94-1, SN5754 47165-04-8, DAPI 47853-44-1,
 SN6113 53222-25-6, SN7167 68772-09-8, SN6999 68772-49-6, SN18071
 88476-80-6, SN 6053 132869-83-1 146426-41-7, SN6131 146426-42-8,
 SN6132 163228-16-8 163228-20-4 213137-22-5
 RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)
 (mol. modeling and footprinting studies of DNA minor groove binders using bisquaternary ammonium heterocyclic compds.)

RE.CNT 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD


RE

- (1) Rydzewski, J; Nucleic Acids Research 1996, V24, P1287 HCAPLUS
- (2) Sayers, E; Biochemistry 1993, V32, P9094 HCAPLUS
- (3) Squire, C; Nucleic Acids Research 1997, V25, P4072 HCAPLUS
- (4) Sriram, M; Annals of the New York Academy of Sciences 1994, V726, P18 HCAPLUS
- (5) Stewart, J; MOPAC Manual, 6th edn 1990
- (6) Suhnel, J; Computer Applications in the Biosciences 1996, V12, P227 MEDLINE
- (7) Teng, M; Nucleic Acids Research 1988, V16, P2671 HCAPLUS
- (8) Treboux, G; Journal of Physical Chemistry 1995, V99, P6417 HCAPLUS
- (9) Wang, A; Crystallographic and Modeling Methods in Molecular Design 1990, P123 HCAPLUS
- (10) Zakrzewska, K; Nucleic Acids Research 1983, V11, P8825 HCAPLUS
- (11) Zasedatelev, A; FEBS Letters 1991, V281, P209 HCAPLUS


IT 213137-22-5
 RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)
 (mol. modeling and footprinting studies of DNA minor groove binders using bisquaternary ammonium heterocyclic compds.)

RN 213137-22-5 HCAPLUS


CN Phenol, 4-[2,5':2',5''-ter-1H-benzimidazol]-2''-yl- (9CI) (CA INDEX NAME)



L28 ANSWER 11 OF 17 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 1996:607536 HCAPLUS
 DN 125:265169
 ED Entered STN: 12 Oct 1996
 TI Isohelicity and Phasing in Drug-DNA Sequence Recognition: Crystal
 Structure of a Tris(benzimidazole)-Oligonucleotide Complex
 AU Clark, George R.; Gray, Emily J.; Neidle, Stephen; Li, Yu-Hua; Leupin,
 Werner
 CS CRC Biomolecular Structure Unit, Institute of Cancer Research,
 Sutton/Surrey, SM2 5NG, UK
 SO Biochemistry (1996), 35(43), 13745-13752
 CODEN: BICHAW; ISSN: 0006-2960
 PB American Chemical Society
 DT Journal
 LA English
 CC 1-6 (Pharmacology)
 AB The crystal structure is reported of a tris(benzimidazole) analog of the
 minor-groove drug Hoechst 33258 bound to the sequence d(CGCAAATTTGCG)2.
 The structure has been refined to an R factor of 17.4% at a resolution of 2.2
 .ANG.. The ligand covers .apprx.71/2 base pairs, including the 5'-AAATTT
 central sequence. This has an exceptionally narrow minor-groove width,
 together with high propeller twists for individual base pairs. The ligand
 has a highly twisted structure, with an overall twist of 50.degree.
 between aromatic rings. All three benzimidazole subunits are in register
 with the DNA, and there is a sym. group of six hydrogen bonds between
 ligand and A.cntdot.T base-pair edges. By contrast, the ligand does not
 show an optimal isohelical fit to the DNA. The correct phasing of drug
 and DNA base pairs is ensured by a number of changes to the DNA such that the
 central 5'-AAATTT region is slightly unwound relative to the structures of
 other noncovalent minor-groove drug complexes.
 ST drug DNA sequence recognition crystal structure; trisbenzimidazole
 oligonucleotide complex crystal structure
 IT Crystal structure
 Hydrogen bond
 (isohelicity and phasing in drug-DNA sequence recognition using crystal
 structure of a benzimidazole-oligonucleotide complex)
 IT Deoxyribonucleic acids
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL
 (Biological study); PROC (Process)
 (isohelicity and phasing in drug-DNA sequence recognition using crystal
 structure of a benzimidazole-oligonucleotide complex)
 IT 149318-33-2 182496-21-5
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL
 (Biological study); PROC (Process)
 (isohelicity and phasing in drug-DNA sequence recognition using crystal
 structure of a benzimidazole-oligonucleotide complex)
 IT 182496-22-6
 RL: FMU (Formation, unclassified); PRP (Properties); FORM (Formation,
 nonpreparative)
 (isohelicity and phasing in drug-DNA sequence recognition using crystal
 structure of a benzimidazole-oligonucleotide complex)
 IT 182496-21-5
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL
 (Biological study); PROC (Process)
 (isohelicity and phasing in drug-DNA sequence recognition using crystal
 structure of a benzimidazole-oligonucleotide complex)
 RN 182496-21-5 HCAPLUS
 CN 3-Pyrrolidinamine, 1-[2''-(4-methoxyphenyl)[2,5':2',5''-ter-1H-
 benzimidazol]-5-yl]- (9CI) (CA INDEX NAME)



IT 182496-22-6
 RL: FMU (Formation, unclassified); PRP (Properties); FORM (Formation, nonpreparative)
 (isohelicity and phasing in drug-DNA sequence recognition using crystal structure of a benzimidazole-oligonucleotide complex)
 RN 182496-22-6 HCAPLUS
 CN Guanosine, 2'-deoxycytidylyl-(5'.fwdarw.3')-2'-deoxyguanylyl-(5'.fwdarw.3')-2'-deoxycytidylyl-(5'.fwdarw.3')-2'-deoxyadenylyl-(5'.fwdarw.3')-2'-deoxyadenylyl-(5'.fwdarw.3')-2'-deoxyadenylyl-(5'.fwdarw.3')-thymidylyl-(5'.fwdarw.3')-thymidylyl-(5'.fwdarw.3')-2'-deoxyguanylyl-(5'.fwdarw.3')-2'-deoxycytidylyl-(5'.fwdarw.3')-2'-deoxy-, double-stranded complementary, compd. with 1-[2'-(4-methoxyphenyl)[2,5':2',5''-ter-1H-benzimidazol]-5-yl]-3-pyrrolidinamine (1:1) (9CI) (CA INDEX NAME)
 CM 1
 CRN 182496-21-5
 CMF C32 H28 N8 O




CM 2
 CRN 146217-99-4
 CMF Unspecified
 CCI MAN

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

L28 ANSWER 12 OF 17 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 1975:112032 HCAPLUS
 DN 82:112032
 ED Entered STN: 12 May 1984
 TI Basic substituted 2,6-bisbenzimidazole derivatives, a novel series of substances with chemotherapeutic activity
 AU Loewe, H.; Urbanietz, J.
 CS Hoechst A.-G., Frankfurt/Main, Fed. Rep. Ger.
 SO Arzneimittel-Forschung (1974), 24(12), 1927-33
 CODEN: ARZNAD; ISSN: 0004-4172
 DT Journal
 LA German
 CC 28-18 (Heterocyclic Compounds (More Than One Hetero Atom))
 GI For diagram(s), see printed CA Issue.
 AB Reaction of 2,5-O₂NClC₆H₃NH₂ with RH [R = R₁ (with R₂ = Me, Et, CHMe₂, Bu, CH₂Ph, CH₂CH₂OH, CO₂Et, CH₂CH₂NEt₂, Ph, CONEt₂, or 2-pyridinyl), piperidino, morpholino, or NEt₂] gave 2,5-O₂NRC₆H₃NH₂, which were reduced to give 3,4-(H₂N)2-C₆H₃R (I). I reacted with 3,4-O₂N(H₂N)C₆H₃C(:NH)OEt.HCl to give the benzimidazoles II (R₃ = NO₂), reduction of which over Raney Ni gave II (R₃ = NH₂), which reacted with 2,3,4-R₆-R₄R₅C₆H₂C(:NH)OEt.HCl to give III (R₄ = H, Cl, Me, NO₂, or OMe; R₅ = H, OMe, OEt, OPr, OBU, Me, Cl, NMe₂, NH₂ OPh, Ph, NO₂, or OH; or R₄R₅ = OCH₂O; R₆ = H or OH). III had anthelmintic activity, especially against filarias in cotton rats. In addition III showed fluorochromic properties.
 ST benzimidazole piperazinylbis anthelmintic; piperazinylbisbenzimidazole

Search done by Noble Jarrell

anthelmintic
 IT Anthelmintics
 (2,6'-bibenzimidazoles as)
 IT 54998-08-2P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
 (Reactant or reagent)
 (preparation and reaction with aminonitrobenzimidoyl ethyl ether)
 IT 23470-57-7P 23470-58-8P 23470-59-9P 23470-60-2P 23470-62-4P
 23491-49-8P 23617-82-5P 23617-83-6P 54998-09-3P 54998-10-6P
 54998-26-4P 54998-27-5P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
 (Reactant or reagent)
 (preparation and reaction with benzimidoyl ethyl ether)
 IT 23470-40-8P 23470-41-9P 23470-42-0P 23470-43-1P 23470-44-2P
 23470-45-3P 23470-47-5P 23470-49-7P 23470-50-0P 23470-51-1P
 23470-52-2P 23470-53-3P 23470-54-4P 23470-56-6P 23491-48-7P
 54997-96-5P 54997-97-6P 54997-98-7P 54997-99-8P 54998-00-4P
 54998-01-5P 54998-02-6P 54998-03-7P 54998-04-8P 54998-05-9P
 54998-06-0P 54998-07-1P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
 (Reactant or reagent)
 (preparation and reduction of)
 IT 23470-26-0P 23470-27-1P 23470-28-2P 23470-29-3P 23470-30-6P
 23470-31-7P 23470-33-9P 23491-44-3P 23491-50-1P 23491-51-2P
 23491-52-3P 23491-53-4P 23491-54-5P 23554-98-5P
 23554-99-6P 23555-00-2P 23555-01-3P 23555-02-4P 23617-78-9P
 23623-06-5P 23623-07-6P 23623-08-7P 23651-51-6P 23651-52-7P
 23685-00-9P 23813-09-4P 54998-11-7P 54998-12-8P 54998-13-9P
 54998-14-0P 54998-15-1P 54998-16-2P 54998-17-3P 54998-18-4P
 54998-19-5P 54998-20-8P 54998-21-9P 54998-22-0P 54998-23-1P
 54998-24-2P 54998-25-3P 55038-64-7P 55038-65-8P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of)
 IT 4278-03-9 40546-41-6 40546-45-0 54998-35-5 54998-36-6 54998-37-7
 54998-38-8 54998-39-9 54998-40-2 54998-41-3 54998-42-4
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (reaction of, with (diaminophenyl)piperazinobenzimidazoles)
 IT 5333-86-8 43002-64-8 51618-01-0 54998-28-6 54998-29-7 54998-30-0
 54998-31-1 54998-32-2 54998-33-3 54998-34-4
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (reaction of, with (dimethylaminophenyl)piperazinobenzimidazoles)
 IT 92-54-6 103-76-4 109-01-3 110-91-8 119-54-0 120-43-4 2759-28-6
 4038-92-0 4318-42-7 5308-25-8 5610-49-1 34803-66-2
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (reaction of, with chloronitroaniline)
 IT 1635-61-6
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (reaction of, with piperazines)
 IT 109-89-7, reactions 110-89-4, reactions
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (with chloronitroaniline)
 IT 23491-53-4P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of)
 RN 23491-53-4 HCAPLUS
 CN 2,5'-Bi-1H-benzimidazole, 2'-(1,3-benzodioxol-5-yl)-5-(4-methyl-1-
 piperazinyl)- (9CI) (CA INDEX NAME)



L28 ANSWER 13 OF 17 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 1969:481418 HCAPLUS
 DN 71:81418
 ED Entered STN: 12 May 1984
 TI Piperazino bisbenzimidazoles
 PA Farbwerke Hoechst A.-G.
 SO Fr., 14 pp.
 CODEN: FRXXAK

DT Patent
 LA French
 IC C07D; A61K
 CC 28 (Heterocyclic Compounds (More Than One Hetero Atom))
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	FR 1519964		19680405		
	DE 1670684			DE	
	GB 1186723			GB	
	US 3538097		19700000	US	
PRAI	DE		19660401		

CLASS

PATENT NO.	CLASS	PATENT FAMILY CLASSIFICATION CODES
------------	-------	------------------------------------

FR 1519964	IC	C07DIC A61K
------------	----	-------------

GI For diagram(s), see printed CA Issue.

AB A variety of methods may be used to prepare the title compds. (I). Thus, heating 50 g. 5-chloro-2-nitroacetanilide, 30 g. 1-methylpiperazine, and 33 g. K₂CO₃ in 50 ml. Me₂NCHO 4 hrs., addition of 300 ml. H₂O, dissoln. of the precipitate in dilute HCl, and reprecipn. by basification gave 54 g. 5-(1-methyl-4-piperazinyl)-2-nitroacetanilide, m. 135.degree., hydrolysis of which gave 44 g. 5-(1-methyl-4-piperazinyl)-2-nitroaniline (II), m. 155.degree.. Hydrogenation of 40 g. II on Ni in 120 ml. MeOH gave 5-(1-methyl-4-piperazinyl)-1,2-diaminobenzene (III). A mixture of crude III with 51 g. 3,4-(O₂N)(H₂N)C₆H₃C(:NH)OEt.HCl (IIa) and 300 ml. HOAc was heated 6 hrs. on a H₂O bath to give 55 g. 2-(3-nitro-4-aminophenyl)-6-(1-methyl-4-piperazinyl)benzimidazole (IV), m. 183-5.degree.. Hydrogenation of 30 g. IV on Ni in 150 ml. N HOAc at 90.degree. gave 18 g. 2-(3,4-diaminophenyl)-6-(1-methyl-4-piperazinyl)benzimidazole (V), m. 268.degree.. V (205 g.) and 160 g. 4-HOC₆H₄C(:NH)OEt.HCl in 1 l. HOAc was heated 2 hrs. under N on a steam bath giving 110 g. 2-[2-(4-hydroxyphenyl)-6-benzimidazolyl]-6-(1-methyl-4-piperazinyl)benzimidazole (I, Ar = HOC₆H₄, R₁ = Me, R₂ = H) (Ia) dihydrate; anhydride m. 235.degree. (decomposition); Ia.3HCl, decomposing 280.degree.; Ia.H₃PO₄, decomposing 315.degree.. Other I (R₁ = Me, R₂ > H) prepared were (Ar, m.p. anhydrous form, and composition of hydrate given): 2-HOC₆H₄, >200.degree., I.H₂O; 4-MeOC₆H₄, 255.degree., I.1.5H₂O; 3-Me-OC₆H₄, 220.degree., I.0.5H₂O; 4-PrOC₆H₄, 286.degree., I.H₂O; 4-EtOC₆H₄, 268.degree., I.1.5H₂O; 4-BuOC₆H₄, 270.degree., I.0.5H₂O.0.5EtOH; 3,4-CH₂O₂C₆H₃, >200.degree., I.2H₂O; Ph, 190.degree. (di-Bz derivative m. 247.degree.), I.H₂O.0.33PrOH; 4-MeC₆H₄, >200.degree., I.H₂O.0.5EtOH; 3-Me-C₆H₄, 236.degree., I.H₂O; 4-ClC₆H₄, >200.degree., I.1.5H₂O; 4-Me₂NC₆H₄, 210.degree., I.2H₂O; 3-chloro-4-methylphenyl, >200.degree., I.H₂O; 4-chloro-3-methylphenyl, 256.degree., -; 3-nitro-4-aminophenyl, 240.degree., I.2H₂O; 4-PhC₆H₄, 310.degree., I.H₂O; 2-naphthyl, 245.degree., I.0.5H₂O; 3-nitro-4-(2-diethylaminoethylamino)phenyl, 294.degree., I.0.5H₂O; and 4-NO₂C₆H₄ (Ib), 210.degree., I.3H₂O. By similar methods, 50 g. 6-chloro-3-nitro-4-acetamidotoluene and 30 g. 1-methylpiperazine gave 44 g. 4-methyl-5-(1-methyl-4-piperazinyl)-2-nitroacetanilide, m. 166.degree., hydrolysis of which gave 4-methyl-5-(1-methyl-4-piperazinyl)-2-nitroaniline (VI), m. 208.degree.. VI and IIa gave 2-(3-nitro-4-aminophenyl)-5-methyl-6-(1-methyl-4-piperazinyl)benzimidazole, m. 280.degree., which was hydrogenated to give 2-(3,4-diaminophenyl)-5-methyl-6-(1-methyl-4-piperazinyl)benzimidazole (VII), m. 155.degree.. VII (11.3 g.) and 8 g. 4-methoxybenzimidino ether hydrochloride in 80 ml. HOAc gave 10.3 g. 2-[2-(4-methoxyphenyl)-6-benzimidazolyl]-5-methyl-6-(1-methyl-4-piperazinyl)benzimidazole (I, Ar = 4-MeOC₆H₄, R₁ = R₂ = Me) sesquihydrate; anhydride m. 196.degree.. Compds. prepared in similar sequences were: 4-chloro-5-(1-methyl-4-piperazinyl)-2-nitroacetanilide, m. 150.degree., 4-chloro-5-(1-methyl-4-piperazinyl)-2-nitroaniline, m. 202.degree., 4-chloro-5-(1-methyl-4-piperazinyl)-1,2-diaminobenzene, 2-(3-nitro-4-aminophenyl)-5-chloro-6-(1-methyl-4-piperazinyl)benzimidazole, m. 258.degree., and 2-[2-(4-methoxyphenyl)-6-benzimidazolyl]-5-chloro-6-(1-methyl-4-piperazinyl)benzimidazole (I, Ar = 4-MeOC₆H₄, R₁ = Me, R₂ = Cl) tetrahydrate; anhydride m. 207.degree.; 5-(1-ethyl-4-piperazinyl)-2-nitroacetanilide, m. 102.degree., 5-(1-ethyl-4-piperazinyl)-2-nitroaniline, m. 125.degree., 5-(1-ethyl-4-piperazinyl)-1,2-diaminobenzene, 2-(3-nitro-4-aminophenyl)-6-(1-ethyl-4-piperazinyl)benzimidazole, m. 188.degree., 2-(3,4-diaminophenyl)-6-(1-methyl-4-piperazinyl)benzimidazole, m. 170.degree., and 2-[2-(4-methoxyphenyl)-6-benzimidazolyl]-6-(1-ethyl-4-piperazinyl)benzimidazole, m. 188.degree. (monohydrate); 5-(1-isopropyl-4-piperazinyl)-2-nitroacetanilide, m. 87.degree., 5-(1-isopropyl-4-piperazinyl)-2-nitroaniline, m. 127.degree., 5-(1-isopropyl-4-piperazinyl)-1,2-diaminobenzene, m. 131.degree.,

2-(3-nitro-4-aminophenyl)-6-(1-isopropyl-4-piperazinyl)benzimidazole, m. 199.degree., 2-(3,4-diaminophenyl)-6-(1-isopropyl-4-piperazinyl)benzimidazole, m. 284.degree., and 2-[2-(4-methoxyphenyl)-6-benzimidazolyl]-6-(1-isopropyl-4-piperazinyl)benzimidazole, m. 214.degree. (dihydrate-hemisopropylate); 5-(1-butyl-4-piperazinyl)-2-nitroacetanilide, m. 89.degree., 5-(1-butyl-4-piperazinyl)-2-nitroaniline, m. 110.degree., 5-(1-butyl-4-piperazinyl)-1,2-diaminobenzene, 2-(3-nitro-4-aminophenyl)-6-(1-butyl-4-piperazinyl)benzimidazole, m. 170.degree., 2-(3,4-diaminophenyl)-6-(1-butyl-4-piperazinyl)benzimidazole, m. 267.degree., and 2-[2-(4-methoxyphenyl)-6-benzimidazolyl]-6-(1-butyl-4-piperazinyl)benzimidazole, m. 260.degree. (sesquihydrate); 5-(1-benzyl-4-piperazinyl)-2-nitroacetanilide, m. 136.degree.; 5-(1-benzyl-4-piperazinyl)-2-nitroaniline, m. 162.degree.; 5-(1-benzyl-4-piperazinyl)-1,2-diaminobenzene, 2-(3-nitro-4-aminophenyl)-6-(1-benzyl-4-piperazinyl)benzimidazole, m. 170.degree., 2-(3,4-diaminophenyl)-6-(1-benzyl-4-piperazinyl)benzimidazole, m. 206.degree., and 2-[2-(4-methoxyphenyl)-6-benzimidazolyl]-6-(1-benzyl-4-piperazinyl)benzimidazole, m. 169.degree. (hemihydrate); 2-[2-(4-methoxyphenyl)-6-benzimidazolyl]-6-(4-piperazinyl)benzimidazole (by hydrogenolysis of the preceding compound), m. 267.degree. (3.5H₂O); 5-[1-(2-hydroxyethyl)-4-piperazinyl]-2-nitroacetanilide, m. 157.degree., 5-[1-(2-hydroxyethyl)-4-piperazinyl]-2-nitroaniline, m. 160.degree., 5-[1-(2-hydroxyethyl)-4-piperazinyl]-1,2-diaminobenzene, 2-(3-nitro-4-aminophenyl)-6-[1-(2-hydroxyethyl)-4-piperazinyl]benzimidazole, m. 120.degree., 2-(3,4-diaminophenyl)-6-[1-(2-hydroxyethyl)-4-piperazinyl]benzimidazole, m. 200.degree.; and 2-[2-(4-methoxyphenyl)-6-benzimidazolyl]-6-[1-(2-hydroxyethyl)-4-piperazinyl]benzimidazole, m. 190.degree. (sesquihydrate); 5-(1-ethoxycarbonyl Hydrogenation of 28 g. Ib in 400 ml. MeOH on Ni gave 18 g. 2-[2-(4-aminophenyl)-6-benzimidazolyl]-6-(1-methyl-4-piperazinyl)benzimidazole, m. 230.degree. (sesquihydrate). I have anthelmintic activity.

ST bibenzimidazoles piperazino; piperazino bibenzimidazoles; anthelmintic bibenzimidazoles


IT 23470-19-1P 23470-20-4P 23470-21-5P 23470-22-6P 23470-23-7P
 23470-24-8P 23470-25-9P 23470-26-0P 23470-27-1P 23470-28-2P
 23470-29-3P 23470-30-6P 23470-31-7P 23470-32-8P 23470-33-9P
 23491-44-3P 23491-45-4P 23491-46-5P 23491-47-6P 23491-48-7P
 23491-49-8P 23491-50-1P 23491-51-2P 23491-52-3P 23491-53-4P
 23491-54-5P 23491-55-6P 23491-56-7P 23554-98-5P 23554-99-6P
 23555-00-2P 23555-01-3P 23555-02-4P 23555-03-5P 23617-77-8P
 23617-78-9P 23623-05-4P 23623-06-5P 23623-07-6P 23623-08-7P
 23651-51-6P 23651-52-7P 23685-00-9P 23813-09-4P

RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of)

IT 23491-53-4P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of)

RN 23491-53-4 HCAPLUS

CN 2,5'-Bi-1H-benzimidazole, 2'-(1,3-benzodioxol-5-yl)-5-(4-methyl-1-piperazinyl)- (9CI) (CA INDEX NAME)



L28 ANSWER 14 OF 17 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1954:71716 HCAPLUS

DN 48:71716

OREF 48:12740f-g

ED Entered STN: 22 Apr 2001

TI Imidazole derivatives. VII. Preparation of sulfonic acids of benzimidazole by baking method

AU Efros, L. S.

SO Zhurnal Obshchei Khimii (1953), 23, 881-2

CODEN: ZOKHA4; ISSN: 0044-460X

DT Journal

LA English

CC 10 (Organic Chemistry)

AB See C.A. 48, 4524c.

IT Blood
 (-coagulation-inhibiting substances)

IT Spectra
 (of benzimidazole derivs. and polybenzimidazoles)

IT 2,5'(or 2,6')-2',5''(or 2',6'')-Terbenzimidazole, 2'',5(or 2'',6)-dimethyl-, trihydrochloride
 2,5'(or 2,6')-2',5''(or 2',6'')-Terbenzimidazole, 2''-methyl-, trihydrochloride
 2,5'(or 2,6')-2',5''(or 2',6'')-Terbenzimidazole, 2''-phenyl-
 2,5'(or 2,6')-2',5''(or 2',6'')-Terbenzimidazole, 2''-phenyl-, dihydrochloride
 5(or 6)-Benzimidazolecarboxylic acid
 5(or 6)-Benzimidazolecarboxylic acid, sulfate
 5(or 6)-Benzimidazolecarboxylic acid, 2-methyl-
 5(or 6)-Benzimidazolecarboxylic acid, 2-phenyl-, hydrochloride
 5(or 6)-Benzimidazolesulfonic acid
 5(or 6)-Benzimidazolesulfonic acid, 2-methyl-
 Benzimidazole, 2-(3,4-diaminophenyl)-
 Benzimidazole, 2-[2-methyl-5(or 6)-benzimidazolyl]-5(or 6)-[5(or 6)-methyl-2-benzimidazolyl]-, trihydrochloride
 Benzimidazole, 2-methyl-, sulfate
 Benzimidazole, 5(or 6)-(2-benzimidazolyl)-2-[2-methyl-5(or 6)-benzimidazolyl]-, trihydrochloride
 Benzimidazole, 5(or 6)-(2-benzimidazolyl)-2-[2-phenyl-5(or 6)-benzimidazolyl]-
 Benzimidazole, 5(or 6)-(2-benzimidazolyl)-2-[2-phenyl-5(or 6)-benzimidazolyl]-, dihydrochloride
 Benzimidazole, 5(or 6)-[5(or 6)-methyl-2-benzimidazolyl]-2-[2-phenyl-5(or 6)-benzimidazolyl]-
 [2,5'(or 2,6')-Bibenzimidazole]-5(or 6)carboxylic acid, 2'-methyl-, dihydrochloride
 [2,5'(or 2,6')-Bibenzimidazole]-5(or 6)carboxylic acid, 2'-phenyl-
 [2,5'(or 2,6')-Bibenzimidazole]-5(or 6)carboxylic acid, 2'-phenyl-, hydrochloride

IT 288-32-4, Imidazole
 (derivs.)

IT 51-17-2, Benzimidazole
 (poly derivs.)


IT 615-15-6, Benzimidazole, 2-methyl- 41292-72-2, 2,5'(or 2,6')-Bibenzimidazole 59695-31-7, 1H-Tetrazole-5-carboxylic acid, 1-phenyl-, potassium salt 66630-70-4, 5(or 6)-Benzimidazolecarboxylic acid, 2-phenyl- 763140-09-6, 2,5'(or 2,6')-Bibenzimidazole, dihydrochloride 763140-22-3, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-phenyl-, dihydrochloride 763140-28-9, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-phenyl- 763140-45-0, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-methyl-, trihydrochloride 763140-62-1, 2,5'(or 2,6')-Bibenzimidazole, 2'-methyl-5(or 6)-[5(or 6)-methyl-2-benzimidazolyl]-, trihydrochloride 763932-97-4, 2,5'(or 2,6')-Bibenzimidazole, 2'-phenyl-, hydrochloride 763932-98-5, 2,5'(or 2,6')-Bibenzimidazole, 2'-phenyl- 763932-99-6, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-methyl-2'-phenyl-, hydrochloride 763933-00-2, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-methyl-2'-phenyl- (preparation of)

IT 1076-38-6, Coumarin, 4-hydroxy-
 (reactions of)

IT 51-17-2, Benzimidazole
 (sulfonated derivs.)


IT 763140-22-3, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-phenyl-, dihydrochloride 763140-28-9, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-phenyl- 763140-45-0, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-methyl-, trihydrochloride 763140-62-1, 2,5'(or 2,6')-Bibenzimidazole, 2'-methyl-5(or 6)-[5(or 6)-methyl-2-benzimidazolyl]-, trihydrochloride (preparation of)

RN 763140-22-3 HCAPLUS
 CN 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-phenyl-, dihydrochloride (SCI) (CA INDEX NAME)




●2 HCl

RN 763140-28-9 HCAPLUS
 CN 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-phenyl- (5CI)
 (CA INDEX NAME)




RN 763140-45-0 HCAPLUS
 CN 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-methyl-,
 trihydrochloride (5CI) (CA INDEX NAME)



●3 HCl

RN 763140-62-1 HCAPLUS
 CN 2,5'(or 2,6')-Bibenzimidazole, 2'-methyl-5(or 6)-[5(or 6)-methyl-2-benzimidazolyl]-, trihydrochloride (5CI) (CA INDEX NAME)



●3 HCl

L28 ANSWER 15 OF 17 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 1954:71715 HCAPLUS
 DN 48:71715
 OREF 48:12740f
 ED Entered STN: 22 Apr 2001
 TI Imidazole derivatives. VI. Synthesis of some polybenzimidazoles
 AU Porai-Koshits, B. A.; Efros, L. S.; Boichinova, E. S.
 SO Zhurnal Obshchei Khimii (1953), 23, 873-9

Search done by Noble Jarrell

CODEN: ZOKHA4; ISSN: 0044-460X

DT Journal

LA English

CC 10 (Organic Chemistry)

AB See C.A. 48, 4523d.

IT Spectra

(of benzimidazole derivs. and polybenzimidazoles)

IT 2,5'(or 2,6')-2',5''(or 2',6'')-Terbenzimidazole, 2'',5(or 2'',6)-dimethyl-, trihydrochloride
 2,5'(or 2,6')-2',5''(or 2',6'')-Terbenzimidazole, 2''-methyl-, trihydrochloride
 2,5'(or 2,6')-2',5''(or 2',6'')-Terbenzimidazole, 2''-phenyl-, dihydrochloride
 2,5'(or 2,6')-2',5''(or 2',6'')-Terbenzimidazole, 2''-phenyl-, dihydrochloride


IT 288-32-4, Imidazole (derivs.)

IT 41292-72-2, 2,5'(or 2,6')-Bibenzimidazole 763140-09-6, 2,5'(or 2,6')-Bibenzimidazole, dihydrochloride 763140-22-3, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-phenyl-, dihydrochloride 763140-28-9, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-phenyl- 763140-45-0, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-methyl-, trihydrochloride 763140-62-1, 2,5'(or 2,6')-Bibenzimidazole, 2'-methyl-5(or 6)-[5(or 6)-methyl-2-benzimidazolyl]-, trihydrochloride 763932-97-4, 2,5'(or 2,6')-Bibenzimidazole, 2'-phenyl-, hydrochloride 763932-98-5, 2,5'(or 2,6')-Bibenzimidazole, 2'-phenyl- 763932-99-6, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-methyl-2'-phenyl-, hydrochloride 763933-00-2, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-methyl-2'-phenyl- (preparation of)

IT 763140-22-3, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-phenyl-, dihydrochloride 763140-28-9, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-phenyl- 763140-45-0, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-methyl-, trihydrochloride 763140-62-1, 2,5'(or 2,6')-Bibenzimidazole, 2'-methyl-5(or 6)-[5(or 6)-methyl-2-benzimidazolyl]-, trihydrochloride (preparation of)

RN 763140-22-3 HCAPLUS


CN 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-phenyl-, dihydrochloride (5CI) (CA INDEX NAME)



● 2 HCl


RN 763140-28-9 HCAPLUS

CN 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-phenyl- (5CI) (CA INDEX NAME)




RN 763140-45-0 HCAPLUS

CN 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-methyl-, trihydrochloride (5CI) (CA INDEX NAME)



●3 HCl

RN 763140-62-1 HCAPLUS
 CN 2,5'(or 2,6')-Bibenzimidazole, 2'-methyl-5(or 6)-[5(or 6)-methyl-2-benzimidazolyl]-, trihydrochloride (5CI) (CA INDEX NAME)



●3 HCl

L28 ANSWER 16 OF 17 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 1954:71714 HCAPLUS
 DN 48:71714
 OREF 48:12739c-i,12740a-f
 ED Entered STN: 22 Apr 2001
 TI 1,2,4-Triazole analogs of histamine
 AU Ainsworth, C.; Jones, R. G.
 CS Lilly Research Labs., Indianapolis, IN
 SO Journal of the American Chemical Society (1953), 75, 4915-18
 CODEN: JACSAT; ISSN: 0002-7863
 DT Journal
 LA Unavailable
 CC 10 (Organic Chemistry)
 OS CASREACT 48:71714
 AB 3-(2-Aminoethyl)-1H-1,2,4-triazole (I) and several of its derivs. have been synthesized. I and, to a lesser degree, its 3-PhCH2NHCH2CH2 (II), 3-Me2CHNHCH2CH2 (III), and 3-AcNHCH2CH2 analogs (IV) exhibited a typical histaminelike activity and were effective orally. H2NCSNHNH2 (102 g.) and 700 cc. dry pyridine treated during 1-2 h. portion-wise with 237 g. .omicron.-C6H4(CO)2NCH2CH2COCl below 0.degree., the mixture let stand overnight, poured with stirring into 2 l. ice water, and the heavy white precipitate washed with 1 l. ice water, 1 l. 50% aqueous AcOH, and again 1 l. ice water yielded 235-50 g. (80-5%) .omicron.-C6H4(CO)2NHCH2CH2CONHNHCSNH2 (V), white needles, m. 238-9.degree. (decomposition) (from AcOH). V (292 g.), 60 g. NaOMe, and 2.5 l. absolute EtOH refluxed overnight, about 2 l. solvent evaporated in vacuo, the residue added with stirring to 2 l. ice water containing 125 cc. concentrated HCl, the mixture let stand, and the solid product washed with 500 cc. H2O, 200 cc. 50% aqueous AcOH, and 200 cc. glacial AcOH gave 140 g. (50%) 3-(2-phthalimidoethyl)-1H-1,2,4-triazole-5-thiol (VI), white needles, m. 295-7.degree. (from AcOH). VI (13.7 g.) suspended in 50 cc. H2O and treated with 10 cc. N2H4.H2O, the mixture let stand overnight at room temperature, and the resulting solid recrystd. from 200 cc. hot H2O yielded 4.5 g. (61%) 3-(2-aminoethyl)-1H-1,2,4-triazole-5-thiol, needles, m. 296-8.degree. (decomposition); the base dissolved in dilute HCl and the solution concentrated to dryness in vacuo gave the HCl salt, needles, m. 270.degree. (precipitated from MeOH with Et2O). VI (27.4 g.), 5.4 g. NaOMe, and 6.2 cc. MeI in 200 cc. EtOH refluxed 2 h., the solvent evaporated in vacuo, the residue extracted with 150 cc. hot EtOH, and the extract cooled deposited 20 g. 3-(2-phthalimidoethyl)-5-methylthio-1H-1,2,4-triazole (VII), dendritic crystals, m. 170-2.degree. (from H2O). VII (5.8 g.) and 3 cc. N2H4.H2O in 50 cc. H2O let stand overnight at room temperature, the mixture evaporated in vacuo,

Search done by Noble Jarrell


the residue extracted with 100 cc. hot C₆H₆, the extract evaporated, the residue dissolved in EtOH, and the solution treated with dry HCl gave 3.2 g. (70%) di-HCl salt (VIII) of 3-(2-aminoethyl)-5-methylthio-1H-1,2,4-triazole, m. 218.degree. (decomposition) (from MeOH-Et₂O). VII (5.8) and 50 cc. 6N HCl refluxed 6 h., the mixture cooled, filtered, and the filtrate evaporated in vacuo gave 76% VIII. VI (2.7 g.) in 100 cc. 10% aqueous AcOH treated 1 h. below 10.degree. with stirring with dry Cl, the resulting white solid filtered off, added directly to 100 cc. concentrated NH₄OH, the solution evaporated on a steam bath overnight, the solid residue slurried with 50 cc. N HCl, and the product recrystd. from H₂O gave 1.3 g. (40%) 3-(2-phthalimidoethyl)-1H-1,2,4-triazole-5-sulfonamide (IX), m. 280-2.degree. (decomposition). IX (3.2 g.), 3 cc. N₂H₄.H₂O, and 50 cc. MeOH refluxed 0.5 h., the solvent removed in vacuo, the residue dissolved in 50 cc. H₂O, the solution treated with 6N HCl, the precipitated phthalhydrazide filtered off, the filtrate evaporated to dryness, the residue treated with 50 cc. N NaOH, again taken to dryness in vacuo, and treated with 50 cc. 6N HCl, the solution evaporated to dryness, the residue extracted with EtOH, and the extract diluted with Et₂O gave 3-(2-aminoethyl)-1H-1,2,4-triazole-5-sulfonamide-HCl, irregular white prisms, m. 170.degree. (decomposition), also obtained by the hydrolysis of IX with 6N HCl. To 100 cc. concentrated HNO₃, 200 cc. H₂O, and 1 g. NaNO₂ was added below 45.degree. with stirring 100 g. VI in small portions, the mixture cooled to 0.degree., cautiously neutralized with saturated aqueous Na₂CO₃, and the precipitate washed with H₂O to give 40 g. (43%) 3-(2-phthalimidoethyl)-1H-1,2,4-triazole (X), needles, m. 215.degree. (from H₂O); HCl salt, m. 245.degree. (from MeOH-Et₂O). VI dissolved in dilute aqueous NaOH, the solution acidified, and the product oxidized similarly with HNO₃ gave no X. VI (1 g.) 3 teaspoonfuls Raney Ni, and 200 cc. EtOH refluxed 4 h., the hot mixture filtered, and the filtrate evaporated to dryness in vacuo gave X, white needles, m. 214-15.degree.. X (40 g.) and 500 cc. 6N HCl refluxed 8 h., the mixture cooled several hrs., filtered, the filtrate evaporated to dryness in vacuo, the residue dissolved in 500 cc. MeOH, and the solution treated with C and diluted with 1 l. Et₂O gave 25-30 g. (84-97%) I.2HCl, decomposed at 251.degree.. I.2HCl (18.5 g.) in 100 cc. absolute EtOH refluxed 1 h. with 10.8 g. NaOMe, the mixture filtered, and the filtrate distilled gave 9 g. (80%) I, b.p. 158-60.degree., m. 83-5.degree.; dipicrate, yellow cubes, m. 190.degree. (from EtOH). I (11.2 g.), and 5.1 g. Me₂CO in 100 cc. EtOH hydrogenated 6 h. over 0.1 g. PtO₂ while heated with an IR lamp, the mixture filtered, the filtrate evaporated in vacuo, the residue in 50 cc. EtOH added to 46 g. picric acid in 300 cc. 95% EtOH, the solution cooled, and the solid deposit recrystd. twice from 300-cc. portions of 95% EtOH gave 45 g. (72%) dipicrate of III, m. 142-4.degree.; the dipicrate suspended in 200 cc. PhNO₂, the mixture extracted with three 100-cc. portions of concentrated HCl, the extract washed with CHCl₃, evaporated in vacuo, and the residue dissolved in MeOH and precipitated with Et₂O gave III.2HCl, m. 186.degree.. I (3.4 g.) and 3.2 g. freshly distilled BzH in 50 cc. EtOH refluxed 2 h., the mixture hydrogenated over PtO₂ at 40 lb. pressure, filtered, and the filtrate treated with 13.8 g. picric acid in 100 cc. hot EtOH gave 70% dipicrate of II, m. 115-17.degree. (from 50% aqueous EtOH), converted to II.2HCl, m. 220.degree.. I.2HCl (5.6 g.), 2.4 g. KOCN, and 2.5 g. NaHCO₃ in 100 cc. H₂O evaporated on the steam bath, the residue extracted with 50 cc. EtOH, and the extract diluted with 500 cc. Et₂O gave 3-(2-ureidoethyl)-1H-1,2,4-triazole, m. 188-90.degree.. I.2HCl (5.5 g.) in 100 cc. 2N NaOH treated with stirring at 0.degree. with 2.8 g. BzCl, the mixture treated after 2 h. with 25 g. ice, adjusted with concentrated HCl to pH 5, and the precipitate washed with aqueous NaHCO₃ and recrystd. from H₂O gave 3.8 g. (55%) 3-(2-benzamidoethyl)-1H-1,2,4-triazole, feathery plates, m. 189-90.degree.. I.2HCl (5.5 g.) in 50 cc. 2N NaOH treated at 0.degree. with 2 cc. AcOH, the solution after 0.5 h. acidified with 6N HCl, evaporated to dryness, the residue extracted with 100 cc. warm absolute EtOH, and the extract diluted with 400 cc. Et₂O gave IV.HCl, white solid, m. 160.degree.. I (1.7 g.) and 2 g. Ac₂O in 50 cc. glacial AcOH heated 3 h. on the steam bath, the mixture diluted with 25 cc. H₂O, let stand 15 min., concentrated to dryness in vacuo, the residue recrystd. from EtOH gave IV, needles, m. 215-16.degree..

IT Spectra

(of benzimidazole derivs. and polybenzimidazoles)


- IT 2,5'(or 2,6')-2',5''(or 2',6'')-Terbenzimidazole, 2'',5(or 2'',6)-dimethyl-, trihydrochloride
 2,5'(or 2,6')-2',5''(or 2',6'')-Terbenzimidazole, 2''-methyl-, trihydrochloride
 2,5'(or 2,6')-2',5''(or 2',6'')-Terbenzimidazole, 2''-phenyl-, trihydrochloride
 2,5'(or 2,6')-2',5''(or 2',6'')-Terbenzimidazole, 2''-phenyl-, dihydrochloride
 5(or 6)-Benzimidazolecarboxylic acid
 5(or 6)-Benzimidazolecarboxylic acid, sulfate
 5(or 6)-Benzimidazolecarboxylic acid, 2-methyl-
 5(or 6)-Benzimidazolecarboxylic acid, 2-phenyl-, hydrochloride

5(or 6)-Benzimidazolesulfonic acid
 5(or 6)-Benzimidazolesulfonic acid, 2-methyl-
 Acetamide, N-2-s-triazol-3-ylethyl-
 Benzamide, N-2-s-triazol-3-ylethyl-
 Phthalimide, N-2-s-triazol-3-ylethyl-
 Phthalimide, N-2-s-triazol-3-ylethyl-, hydrochloride
 Phthalimide, N-[2-(5-mercapto-s-triazol-3-yl)ethyl]-
 Phthalimide, N-[2-(5-sulfamoyl-s-triazol-3-yl)ethyl]-
 Phthalimide, N-[2-[5-(methylthio)-s-triazol-3-yl]ethyl]-
 Semicarbazide, 1-(3-phthalimidopropionyl)-3-thio-
 Urea, (2-s-triazol-3-ylethyl)-
 [2,5'(or 2,6')-Bibenzimidazole]-5(or 6)carboxylic acid, 2'-methyl-,
 dihydrochloride
 [2,5'(or 2,6')-Bibenzimidazole]-5(or 6)carboxylic acid, 2'-phenyl-
 [2,5'(or 2,6')-Bibenzimidazole]-5(or 6)carboxylic acid, 2'-phenyl-,
 hydrochloride
 s-Triazole, 3-(2-acetamidoethyl)-
 s-Triazole, 3-(2-acetamidoethyl)-, hydrochloride
 s-Triazole, 3-(2-aminoethyl)-5-(methylthio)-, dihydrochloride
 s-Triazole, 3-(2-benzamidoethyl)-
 s-Triazole, 3-(2-ureidoethyl)-
 s-Triazole-3-sulfonamide, 5-(2-aminoethyl)-, hydrochloride
 s-Triazole-3-thiol, 5-(2-aminoethyl)-
 s-Triazole-3-thiol, 5-(2-aminoethyl)-, hydrochloride
 s-Triazole-3-thiol, 5-(2-phthalimidomethyl)-
 IT s-Triazole, 3-(2-benzylaminoethyl)-
 s-Triazole, 3-(2-isopropylaminoethyl)-
 (and derivs.)
 IT 2,5'(or 2,6')-2',5'-(or 2',6')-Terbenzimidazole, 5(or
 6)-methyl-2'-phenyl-
 2,5'(or 2,6')-Bibenzimidazole, 2',5(or 2',6)-dimethyl-
 2,5'(or 2,6')-Bibenzimidazole, 2'-methyl-
 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-[5(or 6)-methyl-2-benzimidazolyl]-
 2'-phenyl-
 (and salts)
 IT 7728-75-8, s-Triazole, 3-(2-aminoethyl)-
 (and derivs.)
 IT 85-41-6, Phthalimide 288-32-4, Imidazole
 (derivs.)
 IT 288-88-0, s-Triazole
 (histamine-related compds.)
 IT 51-17-2, Benzimidazole
 (poly derivs., sulfonated derivs.)
 IT 7730-80-5, Acetamide, N-2-s-triazol-3-ylethyl-, hydrochloride 41292-72-2
 , 2,5'(or 2,6')-Bibenzimidazole 66630-70-4, 5(or 6)-
 Benzimidazolecarboxylic acid, 2-phenyl- 763140-09-6, 2,5'(or
 2,6')-Bibenzimidazole, dihydrochloride 763140-22-3, 2,5'(or
 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-phenyl-,
 dihydrochloride 763140-28-9, 2,5'(or 2,6')-Bibenzimidazole, 5(or
 6)-(2-benzimidazolyl)-2'-phenyl- 763140-45-0, 2,5'(or
 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-methyl-,
 trihydrochloride 763140-62-1, 2,5'(or 2,6')-Bibenzimidazole,
 2'-methyl-5(or 6)-[5(or 6)-methyl-2-benzimidazolyl]-, trihydrochloride
 763932-97-4, 2,5'(or 2,6')-Bibenzimidazole, 2'-phenyl-, hydrochloride
 763932-98-5, 2,5'(or 2,6')-Bibenzimidazole, 2'-phenyl- 763932-99-6,
 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-methyl-2'-phenyl-, hydrochloride
 763933-00-2, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-methyl-2'-phenyl-
 (preparation of)
 IT 51-45-6, Histamine
 (triazole analogs of)
 IT 763140-22-3, 2,5'(or 2,6')-Bibenzimidazole, 5(or
 6)-(2-benzimidazolyl)-2'-phenyl-, dihydrochloride 763140-28-9,
 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-phenyl-
 763140-45-0, 2,5'(or 2,6')-Bibenzimidazole, 5(or
 6)-(2-benzimidazolyl)-2'-methyl-, trihydrochloride 763140-62-1,
 2,5'(or 2,6')-Bibenzimidazole, 2'-methyl-5(or 6)-[5(or
 6)-methyl-2-benzimidazolyl]-, trihydrochloride
 (preparation of)
 RN 763140-22-3 HCAPLUS
 CN 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-phenyl-,
 dihydrochloride (5CI) (CA INDEX NAME)




●2 HCl

RN 763140-28-9 HCAPLUS
 CN 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-phenyl- (5CI)
 (CA INDEX NAME)




RN 763140-45-0 HCAPLUS
 CN 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-methyl-,
 trihydrochloride (5CI) (CA INDEX NAME)



●3 HCl

RN 763140-62-1 HCAPLUS
 CN 2,5'(or 2,6')-Bibenzimidazole, 2'-methyl-5(or 6)-[5(or 6)-methyl-2-benzimidazolyl]-, trihydrochloride (5CI) (CA INDEX NAME)



●3 HCl

L28 ANSWER 17 OF 17 HCAPLUS COPYRIGHT 2005 ACS on STN
 AN 1954:25016 HCAPLUS
 DN 48:25016
 OREF 48:4523d-i,4524a-c
 ED Entered STN: 22 Apr 2001
 TI Imidazole derivatives. VI. Synthesis of some polybenzimidazoles
 AU Porai-Koshits, B. A.; Efros, L. S.; Boichinova, E. S.
 CS Lensovet Technol. Inst., Leningrad

Search done by Noble Jarrell

SO Zhurnal Obshchei Khimii (1953), 23, 835-41

CODEN: ZOKHA4; ISSN: 0044-460X

DT Journal

LA Unavailable

CC 10 (Organic Chemistry)

OS CASREACT 48:25016


AB cf. *ibid.* 697. To 1.32 g. 5-methylbenzimidazole in 10 ml. 1:3 H₂SO₄ was added dropwise at 100-3.degree. 2.4 g. chromic acid in 10 ml. H₂SO₄ (1:3) and the mixture chilled after 15 min., yielding 5-benzimidazolecarboxylic acid sulfate, which with NaOAc gave the free acid, m. 300-25.degree. (from H₂O). This (1.62 g.) and 1.08 g. .omicron.-C₆H₄(NH₂)₂ in 10 ml. 20% HCl heated in sealed tube 4 hrs. at 180-200.degree., then neutralized with NH₄OH and filtered, gave 5-(2'-benzimidazolyl)benzimidazole, isolated as the di-HCl salt, m. 362.degree. (from concentrated HCl); the free base could not be purified owing to the formation of gels. Similar oxidation of 2,5-dimethylbenzimidazole gave 70-5% 2-methyl-5-benzimidazolecarboxylic acid (I), m. 301-2.degree. (from H₂O). This with .omicron.-C₆H₄(NH₂)₂ in 20% HCl as above gave after 40 min. at 180-200.degree. 2-methyl-5-(2'-benzimidazolyl)benzimidazole-2HCl, m. 339-40.degree. (from HCl), which with NH₄OH gave the free base (II), m. 340.degree. (from dilute EtOH); this with NH₄OH-AgNO₃ in EtOH gave a flocculent di-Ag salt; the free base yields a dipicrate, m. 282-2.5.degree.. 3,4-(H₂N)2C₆H₃Me (1.22 g.) and 1.76 g. I in 10 ml. 20% HCl heated in a sealed tube 4 hrs. at 180-200.degree. gave 2,5'-dimethyl-5-(2'-benzimidazolyl)benzimidazole, m. high and unsharp, which gave a di-HCl salt, m. above 360.degree. (from 25% HCl); the free base yields a picrate, m. 274.degree.. This oxidized with chromic acid as above gave 2-methyl-5-(5'-carboxy-2'-benzimidazolyl)benzimidazole-2HCl (III), m. about 350.degree. (from 15% HCl); this, decarboxylated by heating with sodalime at 300.degree. gave II (picrate, m. 274.degree.). III with .omicron.-C₆H₄(NH₂)₂ and 15% HCl 4 hrs. at 180-200.degree. gave 75% 2-methyl-5-[2'-benzimidazolyl-5'-(2''-benzimidazolyl)]benzimidazole-3HCl, m. above 360.degree. (from dilute HCl). Similarly condensation with 3,4-(H₂N)2C₆H₃Me gave 85-90% 2,5'-dimethyl-5-[2'-benzimidazolyl-5'-(2''-benzimidazolyl)]benzimidazole-3HCl, m. about 400.degree. (from dilute HCl). 2-Phenyl-5-methylbenzimidazole with chromic acid in aqueous H₂SO₄ gave 2-phenyl-5-benzimidazolecarboxylic acid, isolated as the HCl salt, m. 304-5.degree. (from aqueous HCl). Electrometric titration of this gives 2 pH breaks; at 8.4 and a weak one whose position is unstated. This heated with .omicron.-C₆H₄(NH₂)₂ in 15% HCl in sealed tube 6 hrs. at 180-200.degree. gave 2-phenyl-5-(2'-benzimidazolyl)benzimidazole, m. 308-10.degree. (from dilute EtOH); HCl salt, m. 323-6.degree. (from dilute HCl). Similarly 3,4-(H₂N)2C₆H₃Me gave 2-phenyl-5-(5'-methyl-2'-benzimidazolyl)benzimidazole, m. 329-31.degree. (from dilute EtOH); HCl salt, m. 311-15.degree. (from dilute HCl). This was oxidized as above to 2-phenyl-5-(5'-carboxy-2'-benzimidazolyl)benzimidazole, isolated as the HCl salt, m. 314-19.degree., which, heated with .omicron.-C₆H₄(NH₂)₂ and 10% HCl, gave 2-phenyl-5-[2'-benzimidazolyl-5'-(2''-benzimidazolyl)]benzimidazole, isolated as the di-HCl salt, does not m. 360.degree. (from aqueous HCl). 3,4-(H₂N)2C₆H₃Me gave 2-phenyl-5-[2'-benzimidazolyl-5'-(5''-methyl-2''-benzimidazolyl)]benzimidazole, isolated as the di-HCl salt, does not m. 360.degree.; the free base is insol. in organic solvents except AcOH in which it forms the corresponding salt. Heating 2.25 g. 3,4-(H₂N)2C₆H₃CO₂H.HCl with .omicron.-C₆H₄(NH₂)₂ and 10 ml. 20% HCl in a sealed tube 40 min. at 180-200.degree. gave 0.1 g. 3,4-diaminophenylbenzimidazole, m. 325-30.degree. (from 10% HCl); this reacts with HNO₂ without forming a diazonium salt; in AcOH it gives a green precipitate with phenanthrenequinone. Condensation with HCO₂H or AcOH gave the previously described bis-benzimidazole derivs. (cf. C.A. 44, 1100b). Benzimidazoles have characteristic absorption maximum at 2700-800, dibenzimidazoles at 3100-200, and tribenzimidazoles at 3400-500 A.; even the latter absorb but weakly in the visible, being pale yellow.

IT Spectra

(of benzimidazole derivs. and polybenzimidazoles)


IT 2,5'(or 2,6')-2',5''(or 2',6'')-Terbenzimidazole, 2'',5(or 2'',6)-dimethyl-, trihydrochloride
 2,5'(or 2,6')-2',5''(or 2',6'')-Terbenzimidazole, 2''-methyl-, trihydrochloride
 2,5'(or 2,6')-2',5''(or 2',6'')-Terbenzimidazole, 2''-phenyl-, 2,5'(or 2,6')-2',5''(or 2',6'')-Terbenzimidazole, 2''-phenyl-, dihydrochloride
 5(or 6)-Benzimidazolecarboxylic acid
 5(or 6)-Benzimidazolecarboxylic acid, sulfate
 5(or 6)-Benzimidazolecarboxylic acid, 2-methyl-
 5(or 6)-Benzimidazolecarboxylic acid, 2-phenyl-, hydrochloride
 Benzimidazole, 2-(3,4-diaminophenyl)-

Benzimidazole, 2-[2-methyl-5(or 6)-benzimidazolyl]-5(or 6)-[5(or 6)-methyl-2-benzimidazolyl]-, trihydrochloride
 Benzimidazole, 5(or 6)-(2-benzimidazolyl)-2-[2-methyl-5(or 6)-benzimidazolyl]-, trihydrochloride
 Benzimidazole, 5(or 6)-(2-benzimidazolyl)-2-[2-phenyl-5(or 6)-benzimidazolyl]-
 Benzimidazole, 5(or 6)-(2-benzimidazolyl)-2-[2-phenyl-5(or 6)-benzimidazolyl]-, dihydrochloride
 Benzimidazole, 5(or 6)-[5(or 6)-methyl-2-benzimidazolyl]-2-[2-phenyl-5(or 6)-benzimidazolyl]-
 [2,5'(or 2,6')-Bibenzimidazole]-5(or 6)carboxylic acid, 2'-methyl-, dihydrochloride
 [2,5'(or 2,6')-Bibenzimidazole]-5(or 6)carboxylic acid, 2'-phenyl-
 [2,5'(or 2,6')-Bibenzimidazole]-5(or 6)carboxylic acid, 2'-phenyl-, hydrochloride
 IT 2,5'(or 2,6')-2',5''(or 2',6'')-Terbenzimidazole, 5(or 6)-methyl-2''-phenyl-
 2,5'(or 2,6')-Bibenzimidazole, 2',5(or 2',6)-dimethyl-
 2,5'(or 2,6')-Bibenzimidazole, 2'-methyl-
 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-[5(or 6)-methyl-2-benzimidazolyl]-2'-phenyl- (and salts)
 IT 288-32-4, Imidazole (derivs.)
 IT 51-17-2, Benzimidazole (poly derivs.)
 IT 41292-72-2, 2,5'(or 2,6')-Bibenzimidazole 66630-70-4, 5(or 6)-Benzimidazolecarboxylic acid, 2-phenyl- 763140-09-6, 2,5'(or 2,6')-Bibenzimidazole, dihydrochloride 763140-22-3, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-phenyl-, dihydrochloride 763140-28-9, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-phenyl- 763140-45-0, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-methyl-, trihydrochloride 763140-62-1, 2,5'(or 2,6')-Bibenzimidazole, 2'-methyl-5(or 6)-[5(or 6)-methyl-2-benzimidazolyl]-, trihydrochloride 763932-97-4, 2,5'(or 2,6')-Bibenzimidazole, 2'-phenyl-, hydrochloride 763932-98-5, 2,5'(or 2,6')-Bibenzimidazole, 2'-phenyl- 763932-99-6, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-methyl-2'-phenyl-, hydrochloride 763933-00-2, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-methyl-2'-phenyl- (preparation of)
 IT 51-17-2, Benzimidazole (sulfonated derivs.)
 IT 763140-22-3, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-phenyl-, dihydrochloride 763140-28-9, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-phenyl- 763140-45-0, 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-methyl-, trihydrochloride 763140-62-1, 2,5'(or 2,6')-Bibenzimidazole, 2'-methyl-5(or 6)-[5(or 6)-methyl-2-benzimidazolyl]-, trihydrochloride (preparation of)
 RN 763140-22-3 HCAPLUS
 CN 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-phenyl-, dihydrochloride (SCI) (CA INDEX NAME)




● 2 HCl

RN 763140-28-9 HCAPLUS
 CN 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-phenyl- (SCI)
 (CA INDEX NAME)



RN 763140-45-0 HCAPLUS


CN 2,5'(or 2,6')-Bibenzimidazole, 5(or 6)-(2-benzimidazolyl)-2'-methyl-, trihydrochloride (5CI) (CA INDEX NAME)



●3 HCl

RN 763140-62-1 HCAPLUS

CN 2,5'(or 2,6')-Bibenzimidazole, 2'-methyl-5(or 6)-[5(or 6)-methyl-2-benzimidazolyl]-, trihydrochloride (5CI) (CA INDEX NAME)



●3 HCl

=> b home

FILE 'HOME' ENTERED AT 13:35:13 ON 14 JAN 2005

=>